

Complementary Iterated Floor Words and the Flora Game

Aviezri S. Fraenkel*

Department of Computer Science and Applied Mathematics
Weizmann Institute of Science
Rehovot 76100, Israel

September 6, 2009

Abstract

Let $\varphi = (1 + \sqrt{5})/2$ denote the golden section. We investigate relationships between unbounded iterations of the floor function applied to various combinations of φ and φ^2 . We use them to formulate an algebraic polynomial-time winning strategy for a new 4-pile take-away game *Flora*, which is motivated by partitioning the set of games into subsets *CompGames* and *PrimGames*. We present recursive, arithmetic and word-mapping winning strategies for it. The arithmetic one is based on the Fibonacci numeration system. We further show how to generate the floor words induced by the iterations using word-mappings and and characterize them using the Fibonacci numeration system. We also exhibit an infinite array of such sequences.

Keywords: floor function, integer part function, combinatorics of words, combinatorial game theory, Fibonacci numeration system

Mathematics Subject Classification: 11B75, 11B39, 91A46

1 Introduction

As customary, we denote by $\lfloor x \rfloor$ the integer part of x , commonly known as the *floor function*. It is the largest integer not exceeding x . Let $\varphi = (1 + \sqrt{5})/2$ denote the golden section.

Two topics motivate this work. On the one hand, we wish to study what happens when we keep iterating the floor function with either φ or φ^2 in various ways. Are any interesting relationships between them discernible even after an unbounded number of iterations, or total chaos takes over?

On the other hand, we aim at shedding more light on the class of impartial take-away games. This class appears to be partitioned into two disjoint

*fraenkel@wisdom.weizmann.ac.il <http://www.wisdom.weizmann.ac.il/~fraenkel>

subclasses: those that are easy to generalize to more than one or two piles, and those for which this seems to be very hard. A well-known representative of the former is Nim [2], and of the latter Wythoff's game [5]. Some progress in generalizing Wythoff to multiple piles was recently made. See [9], [24], [23]. Three-pile games that are *extensions* rather than generalizations of Wythoff were also given recently [8], [4].

Here we consider an extension of Wythoff to four piles. The efforts in defining a "right" extension and particularly in proving the validity of the winning strategy are considerably greater than those for three-pile extensions. We present four formulations of the winning strategy.

In §2 we investigate unbounded iterations of the floor function and formulate a wealth of relationships and identities. In §3 we define the subclasses *CompGames* and *PrimGames*, which motivate the definition of the 4-pile game dubbed Flora. In §3.1 we formulate an algebraic winning strategy for the game, based on the results derived in §2, and prove that its complexity is poly-time. In §3.2 we formulate a recursive winning strategy which appears very simple, but its polynomiality is implied only by a recent result [12]. We end in §3.3 with a polynomial-time arithmetic winning strategy, based on the Fibonacci numeration system. In §4 we indicate how to generate sequences induced by iterations of the floor function using word-mappings. We apply it to one of the sequences in §4.1. In §4.2 we present our fourth, word-mapping, formulation of the winning strategy of the Flora game, which is also polynomial-time. In §4.3 we use results from §3.3 and §2 and make minor use of the language of §4, to characterize the representations of general cases of the special sequences playing a major role in the algebraic formulation of the winning strategy of Flora. In §5 we show, by means of an example, how to produce infinite complementary arrays using sequences induced by the iterations. In the final Epilogue we wrap up and indicate natural further directions of research.

Let $a(n) = \lfloor n\varphi \rfloor$, $b(n) = \lfloor n\varphi^2 \rfloor$. It is well-known that the sequences a and b split the positive integers [5], §3. An example of an iterated identity is $a(b(n)) = a(n) + b(n)$. It can be abbreviated as $ab = a + b$, where the product means iteration (composition). We also write a^2 for aa , ab^3a^2 for $abbbaa$, etc. An example of 4 iterated complementary sequences is $a^2 = b - 1$, $ab = a + b$, $ba = a + b - 1$, $b^2 = a + 2b$, since every positive integer is in precisely one of these 4 sequences. We use the notation $w = w_1w_2 \dots w_k$ to denote the word w as well as the (iterated) sequence $w(n)$. If the sequence is intended, we sometimes write $w(n)$ rather than only w . Notice that the product, though not commutative, is associative.

Let $h = b$, $u = a$, and for $k \geq 2$, $h^k = a^{k-1}b$, $u^k = ba^{k-1}$. Let $\Delta a^k(n) = a^k(n+1) - a^k(n)$, $\Delta h^k(n) = h^k(n+1) - h^k(n)$. For technical reasons we put

$$a^0(n) = n, \quad h^0(n) = a(n).$$

Further, let $F_{-1} = 1$, $F_0 = 1$, $F_n = F_{n-1} + F_{n-2}$ ($n \geq 1$) be the Fibonacci sequence.

Notation 1. For $k \geq 0$, $s \in \mathbb{Z}$, let $G_k = \cup_{n=1}^{\infty} a^k(n)$, $H_k = \cup_{n=1}^{\infty} h^k(n)$, $U_k = \cup_{n=1}^{\infty} u^k(n)$, $V_2 = \cup_{n=1}^{\infty} b^2(n)$, $G_k - s = \cup_{n=1}^{\infty} (a^k(n) - s)$ (subtracting s from every element of G_k).

In particular, $G_0 = \mathbb{Z}_{\geq 1}$, and $H_0 = G_1 = U_1$.

Note. In our applications, $s \in \{0, -1, -2, 2\}$, most often 0.

2 Identities

After multiplying by the irrational φ and then throwing out the fractional part for an unbounded number of times, one might expect complete chaos among relationships involving a^k , h^k , u^k and b^k . It is thus surprising that there are many striking identities and relationships among them. Our purpose in this section is to prove a selection of them.

Theorem 1. For every $k \in \mathbb{Z}_{\geq 1}$ and every $n \in \mathbb{Z}_{\geq 1}$ the following holds:

- (a) The $k+1$ sequences $G_k, H_k, H_{k-1}, \dots, H_2, H_1$ partition $\mathbb{Z}_{\geq 1}$.
- (b) $u^{k+1} = a^k + a^{k+1} = a^{k+2} + 1$.
- (c) $h^k = a^{k+1} + F_{k-1}$.
- (d) $u^{k+1} = h^{k+1} - F_k + 1 = a^{k+2} + 1$.
- (e) (e1) $h^{k+1} - h^k = a^k + F_{k-2} - 1$.
(e2) $h^{k+1} - a^{k+1} = a^k + F_k - 1$.
(e3) $au^{k+1} = u^{k+2} + 1$.
- (f) (f1) Let

$$S_1 = \{n \in \mathbb{Z}_{\geq 1} : \Delta a(n) = F_0\}, \quad S_2 = \{n \in \mathbb{Z}_{\geq 1} : \Delta a(n) = F_1\}.$$

Then S_1 and S_2 split $\mathbb{Z}_{\geq 1}$; and each of S_1 and S_2 is an infinite set.

- (f2) For all $k \in \mathbb{Z}_{\geq 1}$ the following holds: $\Delta a^k(n) = F_{k-1}$ for all $n \in S_1$ and $\Delta a^k(n) = F_k$ for all $n \in S_2$.
- (f3) (i) If $\Delta a^k(n+1) = F_{k-1}$ for some $n \in \mathbb{Z}_{\geq 1}$, then $\Delta a^k(n) = \Delta a^k(n+2) = F_k$. (ii) If $\Delta a^k(n+1) = \Delta a^k(n+2) = F_k$ for some $n \in \mathbb{Z}_{\geq 1}$, then $\Delta a^k(n) = \Delta a^k(n+3) = F_{k-1}$.
- (f4) $\Delta a^k(n) = \Delta h^{k-1}(n) \in \{F_{k-1}, F_k\}$, and each of F_{k-1} and F_k is assumed for infinitely many n .
- (f5) (i) $\Delta a^k(0) = 1$, (ii) $\Delta h^k(0) = F_{k-1} + 1$.
- (g) $a^k(h(n)) = h^{k+1}(n)$ (due to Lior Goldberg).
- (h) (h1) $(G_2 + 2) \subset G_1$.

$$\text{(h2)} \quad G_2 \cup (G_2 + 2) = G_1.$$

$$\text{(h3)} \quad U_2 \subset (G_1 - 2) \subset G_1 \cup U_2.$$

$$\text{(h4)} \quad (V_2 - 1) \subset G_2.$$

We begin by recalling some elementary properties of the floor function. Let x, y be any real numbers. Denote by $\{x\}$ the fractional part of x , so $x = \lfloor x \rfloor + \{x\}$. Then:

- $0 \leq \{x\} < 1, x - 1 < \lfloor x \rfloor \leq x$. Replacing x by $-x$, $-x - 1 < \lfloor -x \rfloor \leq -x$, hence $-1 \leq \lfloor x \rfloor + \lfloor -x \rfloor \leq 0$ and $\lfloor x \rfloor + \lfloor -x \rfloor = 0$ if and only if x is an integer. For example, $\lfloor \varphi \rfloor = 1, \lfloor -\varphi \rfloor = -2, \lfloor \varphi \rfloor + \lfloor -\varphi \rfloor = -1$; and $\varphi^2 = \varphi + 1$ implies $\{ \varphi \} = \varphi^{-1} = \varphi - 1$.
- $\lfloor x \rfloor + \lfloor y \rfloor \leq \lfloor x + y \rfloor \leq \lfloor x \rfloor + \lfloor y \rfloor + 1$. This follows immediately from $\lfloor x + y \rfloor = \lfloor \lfloor x \rfloor + \{x\} + \lfloor y \rfloor + \{y\} \rfloor = \lfloor x \rfloor + \lfloor y \rfloor + \lfloor \{x\} + \{y\} \rfloor$.

Lemma 1. (i) Let $s \in \mathbb{Z}$. Each of the sequences $G_k + s, H_k + s, U_k + s, V_2 + s$ is strictly increasing for every $k \geq 1$.

(ii) The sequences G_k, H_k split G_{k-1} for every $k \geq 1$.

Proof. (i) Follows from the fact that $\varphi^2 = \varphi + 1 > \varphi > 1$.

(ii) Since $\varphi^{-1} + \varphi^{-2} = 1$, the sequences G_1 and H_1 split $\mathbb{Z}_{\geq 1} = G_0$ (see e.g., [5], §3), so the result holds for $k = 1$. For any $k \geq 1$, assume that G_k, H_k split G_{k-1} . Then

$$\begin{aligned} G_{k+1} \cup H_{k+1} &= \bigcup_n (a^{k+1}(n) \cup h^{k+1}(n)) = \bigcup_n (aa^k(n) \cup ah^k(n)) \\ &= \bigcup_n aa^{k-1}(n) \text{ (by induction)} = \bigcup_n a^k(n) = G_k. \quad \blacksquare \end{aligned}$$

Proof of Theorem 1 (a). We noted that G_1 and H_1 split $\mathbb{Z}_{\geq 1}$. Suppose that $G_k, H_k, H_{k-1}, \dots, H_2, H_1$ partition $\mathbb{Z}_{\geq 1}$. Then $G_{k+1}, H_{k+1}, H_k, \dots, H_2, H_1$ partition $\mathbb{Z}_{\geq 1}$, since G_{k+1}, H_{k+1} split G_k by Lemma 1. \blacksquare

Note. It follows from Lemma 1(ii) (or from (a)) that for any positive integers m, n , $a(m) \neq b(n)$. This property will be referred to in the sequel as *disjointness*.

Proof of Theorem 1 (b). By definition,

$$a^{k+2} = a^2 a^k = \lfloor \varphi \lfloor \varphi a^k \rfloor \rfloor \leq \lfloor \varphi^2 a^k \rfloor = ba^k = u^{k+1}.$$

By disjointness, $u^{k+1} = ba^k \geq a^{k+2} + 1$. Conversely, multiply $\varphi a^k < a^{k+1} + 1$ by φ to get, $\varphi^2 a^k < \varphi(a^{k+1} + 1)$, hence $ba^k \leq \lfloor \varphi(a^{k+1} + 1) \rfloor$. By disjointness this inequality is strict, so

$$u^{k+1} \leq \lfloor \varphi(a^{k+1} + 1) \rfloor - 1 \leq a^{k+2} + \lfloor \varphi \rfloor = a^{k+2} + 1.$$

On the other hand, $u^{k+1} = \lfloor (\varphi + 1)a^k \rfloor = a^k + a^{k+1}$. \blacksquare

Lemma 2. For every $k \in \mathbb{Z}_{\geq 1}$,

- (i) $\lfloor \varphi F_{k-1} \rfloor \in \{F_k - 1, F_k\}$.
- (ii) $\lfloor \varphi^2 F_{2k-2} \rfloor = F_{2k} - 1$, $\lfloor F_{2k-1} \varphi^2 \rfloor = F_{2k+1}$.

Proof. (i) The ratios F_k/F_{k-1} are the convergents of the simple continued fraction expansion of $\varphi = [1, 1, 1, \dots]$. Therefore $|\varphi F_{k-1} - F_k| < F_{k-1}^{-1}$ (see e.g., [14], ch. 10), so $\varphi F_{k-1} - F_k = \delta$, where $-F_{k-1}^{-1} < \delta < F_{k-1}^{-1}$. Thus $\lfloor \varphi F_{k-1} \rfloor = F_k + \lfloor \delta \rfloor$. The result follows if $|\delta| < 1$, which is the case for all $k \geq 1$, since $F_{k-1} \geq F_0 = 1$.

(ii) The ratios F_{k+2}/F_k are the convergents of the simple continued fraction expansion of $\varphi^2 = [2, 1, 1, 1, \dots]$. In fact, $F_{2k+1}/F_{2k-1} < \varphi^2 < F_{2k}/F_{2k-2}$. This follows easily from [14], ch. 10. Then $\varphi^2 F_{2k-1} - F_{2k+1} = \delta$, where $0 < \delta < F_{2k-1}^{-1}$, hence $\lfloor \varphi^2 F_{2k-1} \rfloor = F_{2k+1}$, since $0 < \delta < 1$ for all $k \geq 1$. Similarly, $\varphi^2 F_{2k-2} - F_{2k} = \delta$, where $-F_{2k-2}^{-1} < \delta < 0$. Thus $\lfloor \varphi^2 F_{2k-2} \rfloor = F_{2k} - 1$, since $-1 < \delta < 0$ for all $k \geq 1$. ■

Lemma 3. $h^2 - a^3 = 2$.

Proof. In Lemma 9 of [8] we proved the special case $k = 1$ of **(d)**, namely $h^2 = u^2 + 1$. Thus $h^2 - a^3 = u^2 - a^3 + 1$. Clearly $a^3 = \lfloor \varphi \lfloor \varphi \lfloor n\varphi \rfloor \rfloor \rfloor \leq \lfloor \varphi^2 a \rfloor = u^2$. But this inequality is strict by disjointness. Thus $h^2 - a^3 \geq 2$.

Conversely, multiply the inequality $\varphi a < a^2 + 1$ by φ , to get $\varphi^2 a < \varphi(a^2 + 1)$. Therefore $\lfloor \varphi^2 a \rfloor \leq \lfloor \varphi(a^2 + 1) \rfloor$. Again by disjointness, this inequality is strict, i.e., $u^2 \leq a^3 + 1$. As we saw, Lemma 9 of [8] asserts $u^2 = h^2 - 1$. Therefore, $h^2 - a^3 \leq 2$. ■

Notation 2. For any positive integer N , denote by $R(N)$ the representation of N in the Fibonacci numeration system. It has the form $R(N) = (d_m, \dots, d_0)$ where $N = \sum_{i=0}^m d_i F_i$, $d_i \in \{0, 1\}$, $d_i = 1 \implies d_{i-1} = 0$, $i \geq 1$ [6]. The *position* of a representation is the subscript i of d_i . Thus, d_0 is in position 0, d_1 in position 1, etc.

Proof of Theorem 1 (c). For $k = 1$, this is Lemma 5 of [8], Sect. 5. For $k = 2$, it is Lemma 3 above. Suppose that $h^k = a^{k+1} + F_{k-1}$ for some arbitrary $k \geq 2$. Multiply by φ and take the floor of both sides. This gives, by Lemma 2, $h^{k+1} = \lfloor \varphi(a^{k+1} + F_{k-1}) \rfloor \leq a^{k+2} + \lfloor \varphi F_{k-1} \rfloor + 1 \leq a^{k+2} + F_k + 1$. Now [8] Sect. 6 implies that $R(a^2)$ ends in 01. By Lemma 1, the same holds for a^k and h^k for every $k \geq 3$ (but it does not hold for h^2). Since $R(F_k)$ ends in 00 for $k \geq 2$, $R(a^{k+2} + F_k)$ also ends in 01 for $k \geq 2$, and so does h^{k+1} for $k \geq 2$. But $R(a^{k+2} + F_k + 1)$ ends in 10. Hence $h^{k+1} = a^{k+2} + F_k$. ■

Proof of Theorem 1 (d). From **(b)** and **(c)**, $u^{k+1} = a^{k+2} + 1 = h^{k+1} - F_k + 1$. The second follows once more from **(c)**. ■

We note that inspection shows that **(d)** does not hold for $k < 1$.

Proof of Theorem 1 (e1). Subtracting **(c)** from **(c)** with k replaced by $k + 1$ gives $h^{k+1} - h^k = a^{k+2} - a^{k+1} + F_{k-2}$. Substituting the value of a^{k+2}

from **(b)** yields the desired result. ■

(e2) Follows from **(e1)**, where we replace h^k by its value from **(c)**. ■

(e3) We have,

$$\begin{aligned} au^{k+1} &= \lfloor \varphi \lfloor \varphi^2 a^k \rfloor \rfloor = \lfloor \varphi(a^k + a^{k+1}) \rfloor \\ &\leq a^{k+1} + a^{k+2} + 1 = u^{k+2} + 1, \end{aligned}$$

where the last equality follows from **(b)**. On the other hand,

$$\lfloor \varphi(a^k + a^{k+1}) \rfloor \geq a^{k+1} + a^{k+2} = u^{k+2}.$$

Hence by disjointness, $\lfloor \varphi u^{k+1} \rfloor = u^{k+2} + 1$. ■

We recall the following special case of Lemma 2 of [5]:

Lemma I. *For integers $i > j \geq 0$ and integer $N_{i+1} \in \mathbb{Z}_{\geq 1}$, let $R(N_{i+1}) = F_i + F_{i-2} + \dots + F_j$, where $j = 0$ if i is even, $j = 1$ if i is odd. Then $N_{i+1} = F_{i+1} - 1$.*

This is the analog in the Fibonacci numeration system of the decimal $99 \dots 9$.

Proof of Theorem 1 (f1). For any $n \in \mathbb{Z}_{\geq 1}$, clearly $\varphi - 1 < \Delta a(n) < \varphi + 1$, so $\Delta a(n) \in \{1, 2\} = \{F_0, F_1\}$. This shows already that S_1, S_2 split $\mathbb{Z}_{\geq 1}$. Moreover, if $\Delta a(n) = 1$ for all large n , then, since $h(n)$ is increasing, we would have $a(n) \cap h(n) \neq \emptyset$ for infinitely many $n \in \mathbb{Z}_{\geq 1}$, contradicting the complementarity of the 2 sequences. If $\Delta a(n) = 2$ for all large n , then also $\Delta h(n) = 2$ for all large n by complementarity. But a direct computation shows that $\Delta a(n) = 2 \implies \Delta h(n) = 3$, another contradiction. Thus each of S_1 and S_2 is infinite as claimed. ■

(f2) We proceed by induction on k . Suppose that for some $k \geq 1$, $\Delta a^k(n) = F_{k-1}$ for all $n \in S_1$, and $\Delta a^k(n) = F_k$ for all $n \in S_2$. This holds for $k = 1$ by **(f1)**. For now let's assume that $n \in S_1$. Then

$$\Delta a^{k+1}(n) = \lfloor \varphi a^k(n+1) \rfloor - \lfloor \varphi a^k(n) \rfloor < \varphi a^k(n+1) - \varphi a^k(n) + 1 = \varphi F_{k-1} + 1$$

by the induction hypothesis. Also,

$$\Delta a^{k+1}(n) > \varphi a^k(n+1) - \varphi a^k(n) - 1 = \varphi F_{k-1} - 1.$$

So $\lfloor \varphi F_{k-1} \rfloor \leq \Delta a^{k+1}(n) \leq \lfloor \varphi F_{k-1} \rfloor + 1$. Then Lemma 2 implies that $\Delta a^k(n) \in \{F_k - 1, F_k, F_k + 1\}$.

In the proof of **(c)** above, it was mentioned that $R(a^2(n+2))$ ends in 01. The same thus holds for $R(a^{k+1}(n+1))$ and $R(a^{k+1}(n))$ for all $k \geq 1$, since G_{k+1} is a subsequence of G_2 for all $k \geq 1$. Therefore $R(\Delta a^{k+1}(n))$ ends in 00, the same as $R(F_k)$. But $R(F_k + 1)$ ends in 01, and Lemma I implies that $R(F_k - 1)$ ends in 10, or in 01, depending on whether k is even or odd. Hence $\Delta a^{k+1}(n) = F_k$ for all $n \in S_1$. The same proof shows that $\Delta a^{k+1}(n) = F_{k+1}$ for all $n \in S_2$. ■

(f3) This follows easily for $k = 1$ by considering the size of φ . For all $k \geq 1$ it follows from **(f2)**. ■

(f4). This follows directly from **(f1)** and **(f2)**. ■

Note. Part of the proof of **(f4)** follows directly from **(d)**:

$$h^{k+1}(n+1) - a^{k+2}(n+1) = h^{k+1}(n) - a^{k+2}(n) = F_k.$$

Hence $\Delta h^{k+1}(n) = \Delta a^{k+2}(n)$. But this establishes the equality part of **(f4)** only for $k \geq 3$ and doesn't prove the membership part.

(f5). (i) Induction on k . (ii) By definition, $\Delta h^k(0) = h^k(1) - h^k(0) = h^k(1)$. The result for $h^k(1)$ follows directly from (i) and **(c)**. ■

(g). For $k = 1$, $h^2(n) = \lfloor \varphi h(n) \rfloor = a(h(n))$. If the assertion holds for any $k \geq 1$, then $h^{k+2}(n) = \lfloor \varphi h^{k+1}(n) \rfloor = \lfloor \varphi a^k(h(n)) \rfloor = a^{k+1}(h(n))$. ■

(h1). Clearly $G_2 \subset G_1$, so for every $n \in \mathbb{Z}_{\geq 1}$, $a^2(n) = a(m)$ for some $m \in \mathbb{Z}_{\geq 1}$. By **(c)**, $a^2(n) + 1 = h(n) \notin G_1$. But then $a^2(n) + 2 = a(m+1) \in G_1$ by **(f4)** for $k = 1$. ■

The following is a special case of *Property 1*, Sect. 5 of [5]:

Lemma II. *The set of numbers $\{R(N) : N \in G_1\}$ ends in an even (possibly 0) number of 0s, hence the complementary set of numbers $\{R(N) : N \in H_1\}$ ends in an odd number of 0s.*

(h2). By **(h1)**, $G_2 \cup (G_2 + 2) \subseteq G_1$. Choose any $a(n) \in G_1$. If $a(n) \in (G_2 + 2)$, we are done. So suppose that $a(n) = a^2(m) + 2$ for no $m \in \mathbb{Z}_{\geq 1}$. By **(c)**, $a^2(m) + 1 = h(m)$ for all $m \in \mathbb{Z}_{\geq 1}$, so by disjointness, $a(n) = a^2(m) + 1$ for no $m \in \mathbb{Z}_{\geq 1}$. But then $a(n) = g^2(m)$ for some $m \in \mathbb{Z}_{\geq 1}$ by **(f4)** for $k = 1$, so $a(n) \in G_2$. ■

(h3). The following is immediately implied by **(f3)**: (a) if $a(n) - 1 \notin G_1$, then $a(n) - 2 \in G_1$; and, conversely, (b) if $a(n) - 1 \in G_1$, then $a(n) - 2 \notin G_1$. Consider case (b). Lemma II then implies that $R(a(n) - 1)$ ends in an even positive number of 0s, and $R(a(n))$ ends in 01. By Lemma I, $R(a(n) - 2)$ then ends in 10. We now show that $R(\lfloor \varphi^2 a(n) \rfloor)$ ends in 10 for all $n \in \mathbb{Z}_{\geq 1}$.

Now $R(a(n))$ ends in F_{2k-2} for some $k \in \mathbb{Z}_{\geq 1}$. By Lemma 2(ii), $\lfloor \varphi^2 F_{2k-2} \rfloor = F_{2k} - 1$, and $R(F_{2k} - 1)$ ends in 10 by Lemma I, the same as $R(a(n) - 2)$ for case (b). This proves that $R(\lfloor \varphi^2 a(n) \rfloor)$ ends in 10 for all $n \in \mathbb{Z}_{\geq 1}$, and the right-hand-side of **(h3)**. On the other hand, let $N \in U_2$. Then $R(N)$ ends in 10, and so $N + 1$ and $N + 2$ are both in G_1 . Thus $N \in G_1 - 2$, proving the left-hand-side of **(h3)**. ■

(h4). In the proof of **(h3)** we showed that $R(\lfloor \varphi^2 a(n) \rfloor)$ ends in 10 for all $n \in \mathbb{Z}_{\geq 1}$. Since $R(h(n))$ ends in an odd number of 1s for all $n \in \mathbb{Z}_{\geq 1}$ by Lemma II, $R(v^2(n))$ ends in an odd number $N \geq 3$ of 1's. Then Lemma I implies that $R(v^2(n) - 1)$ ends in 01. Theorem 3 of [8] states that $R(G_2)$ is the set of all numbers whose representation ends in 01, so $(V_2 - 1) \subset G_2$. ■

Remark. Consider the word $w = \ell_1 \ell_2 \dots \ell_k$ of length k over the binary alphabet $\{a, b\}$. The number m of occurrences of the letter b is the *weight* of w . We also put $F_{-2} = 0$. Recently, Clark Kimberling [18] proved the following nice and elegant result:

Theorem I. For $k \geq 2$, let $w = \ell_1 \ell_2 \dots \ell_k$ of length k be any word over $\{a, b\}$ of length k and weight m . Then $w = F_{k+m-4}a + F_{k+m-3}b - c$, where $c = F_{k+m-1} - w(1) \geq 0$ is independent of n .

Notice that in the theorem only the weight m appears, not the locations within w where the b s appear. The locations, however, obviously influence the behavior of w . This influence is hidden in the “constant” $c = c_{k,m}$, where not only F_{k+m-1} obviously depends on k and m , but also $w(1)$.

We could have used Theorem I to prove most of the results of Theorem 1 simply by expressing each side of an identity as in Theorem I and verifying that they are identical. This verification, however, seems less satisfactory than the above proofs, which shed some light on the nature of the identities. In a recent book review it says, “but it is fair to say that while it is a proof, it is not an explanation” (AMM, **116**, Aug-Sept 2009, p. 660). Hardy [13], writing about seven proofs of the Rogers-Ramanujan identities, put it this way: “None of these proofs can be called “simple” and “straightforward”, since the simplest are essentially verifications”. I got the Hardy reference from opinion 90 on the webpage of my esteemed opinionated friend Doron Zeilberger <http://www.math.rutgers.edu/~zeilberg/OPINIONS.html>. Moreover, the computation of c is not, generally, so easy, as acknowledged by Kimberling. For example, we can show that for $w = h^k$ ($m = 1$) we get $w(1) = h^k(1) = F_{k-1} + 1$, so $c_k = F_{k-2} - 1$. The proof depends on Lemma 2 and the Fibonacci numeration system.

3 An Application: The Flora Game

The class of impartial take-away games appears to be partitioned into two disjoint subclasses:

- *CompGames* (composite games), and
- *PrimGames* (prime games).¹

Informally, *CompGames* are games that are easy to generalize to more than one or two piles; *PrimGames* are those for which this seems to be very hard. A well-known representative of the former is Nim, and of the latter, Wythoff’s game. Some progress in generalizing Wythoff to multiple piles was recently made. Two 3-pile games that are *extensions* rather than generalizations of Wythoff were also given recently. It appears that, largely, a game belongs to class *CompGames* if it decomposes into a *disjunctive* sum of subgames, such as Nim, which is the Nim-sum of its pile sizes; and it belongs to class *PrimGames* if it is not decomposable. Hence the names *CompGames* (*composite* games) and *PrimGames* (not decomposable – *prime*). Whereas for the former there are theories for both the impartial as well as for the partizan case, there is no general theory for the latter yet, and we believe that these “lone wolf” games should be investigated more seriously.

¹It’s different from the partition into *MathGames* and *PlayGames* defined in [7].

Here we study an extension of Wythoff to *four* piles, which appears to be a PrimGame. The efforts in defining a “right” extension, and particularly in proving the validity of the winning strategy, are considerably greater than those for *three*-pile extensions. We present four winning strategies, algebraic, recursive, arithmetic and word-mapping. The recursive is the easiest to describe, though it seems to be hardest computationally. Actually it’s also polynomial-time [12]. The algebraic depends on iterations of the floor function, the arithmetic on the Fibonacci numeration system and the word-mapping on a morphism-like mapping. All are polynomial time winning strategies.

The Flora game is a 2-player game played on 4 piles of tokens. We denote positions of Flora by (a_1, a_2, a_3, a_4) with $0 \leq a_1 \leq a_2 \leq a_3 \leq a_4$. It goes without saying that every pile must contain a nonnegative number of tokens at all times.

The end position is $T_0 := (0, 0, 0, 0)$. The first player unable to move (because the present position is T_0) loses; the opponent wins.

There are 3 rules of move:

I. Any positive number of tokens from up to 3 piles may be removed.

II. From a nonzero position one can move to T_0 if any of the following 3 conditions hold: (i) two piles have the same size, (ii) $a_3 - a_2 = 1$, (iii) $a_1 = h(n)$ and $a_2 < h^2(n) - 2$ for some $n \in \mathbb{Z}_{\geq 1}$.

III. If $0 < a_1 < a_2 < a_3 < a_4$, one can remove $p > 0$ from a_3 , $q > 0$ from a_4 and an arbitrary nonnegative integer from a_1 and a_2 , subject to:

(i) $q = p$ if $a_4 - a_3 \notin V_2$, except for the proviso that if $a_3 - p$ is the second smallest component in the quadruple moved to, then $p \neq 5$.

(ii) $q = p + 1$ if $a_4 - a_3 \in V_2$.

We say that a move in Flora is *legal* if it is consistent with the rules (I)-(III).

Note. If the position moved to under rule III(i) is (b_1, b_2, b_3, b_4) (where of course $0 \leq b_1 \leq b_2 \leq b_3 \leq b_4$), then $a_3 - p = b_i$, $a_4 - p = b_j$ for some $1 \leq i < j \leq 4$. Then $a_4 - a_3 = b_j - b_i = t$ for some $t \in \mathbb{Z}_{\geq 1}$, and normally $t \neq p$.

3.1 Algebraic Formulation of the P -positions

The set of P -positions of a game is the set of game positions from which the second (*Previous*) player can force a win. The set of all P -positions of a game is denoted \mathcal{P} . In particular, for Flora, $T_0 \in \mathcal{P}$.

Let

$$\begin{aligned} A_n &= h(n), & B_n &= a^3(n), & C_n &= h^2(n), & D_n &= h^3(n), \\ A &= \cup_{n=1}^{\infty} A_n, & B &= \cup_{n=1}^{\infty} B_n, & C &= \cup_{n=1}^{\infty} C_n, & D &= \cup_{n=1}^{\infty} D_n, \\ T_n &:= (A_n, B_n, C_n, D_n), & T &= \cup_{n=0}^{\infty} T_n. \end{aligned}$$

A prefix of T of size 19 is shown in Table 1. We shall presently show that T constitutes the set of P -positions of Flora. Assuming the truth of this assertion, we illustrate simple moves in instances of Flora.

- Examples.** (i) From each of the positions $(4, 6, 9, 9)$, $(5, 8, 9, 14)$, $(7, 8, 11, 20)$ one can move to $T_0 \in \mathcal{P}$ by II.
(ii) From $(6, 7, 9, 14)$, one can move to $(5, 6, 8, 12) \in \mathcal{P}$ by I.
(iii) From $(19, 21, 22, 32)$ can move as follows: $19 \rightarrow 9$, $21 \rightarrow 11$, $22 \rightarrow 7$, $32 \rightarrow 17$, resulting in $(7, 9, 11, 17) \in \mathcal{P}$ by III(i).
(iv) From $(24, 29, 32, 37)$, one can move to $(5, 6, 8, 12) \in \mathcal{P}$ by III(ii) (since $37 - 32 = 5 \in V_2$).

Table 1: P -positions of Flora.

n	$h(n)$	$a^3(n)$	$h^2(n)$	$h^3(n)$
0	0	0	0	0
1	2	1	3	4
2	5	6	8	12
3	7	9	11	17
4	10	14	16	25
5	13	19	21	33
6	15	22	24	38
7	18	27	29	46
8	20	30	32	51
9	23	35	37	59
10	26	40	42	67
11	28	43	45	72
12	31	48	50	80
13	34	53	55	88
14	36	56	58	93
15	39	61	63	101
16	41	64	66	106
17	44	69	71	114
18	47	74	76	122

Notation 3. For $n \in \mathbb{Z}_{\geq 1}$, let $\Delta_{DC}(n) := h^3(n) - h^2(n)$, $\Delta_{DB}(n) := h^3(n) - a^3(n)$, $\Delta_{DA}(n) := h^3(n) - h(n)$, $\Delta_{CB}(n) := h^2(n) - a^3(n)$, $\Delta_{CA}(n) := h^2(n) - h(n)$, $\Delta_{BA}(n) := a^3(n) - h(n)$, $\Delta(n) = \Delta_{DC}(n) \cup \Delta_{DB}(n) \cup \Delta_{DA}(n) \cup \Delta_{CB}(n) \cup \Delta_{CA}(n) \cup \Delta_{BA}(n)$, $\Delta = \cup_{n=1}^{\infty} \Delta(n)$.

- Lemma 4.** (i) $\Delta_{DC}(n) = a^2(n)$,
(ii) $\Delta_{DB}(n) = a^2(n) + 2$,
(iii) $\Delta_{DA}(n) = u^2(n)$,
(iv) $\Delta_{CB}(n) = 2$,
(v) $\Delta_{CA}(n) = a(n)$,
(vi) $\Delta_{BA}(n) = a(n) - 2$,
(vii) $\Delta = \mathbb{Z}_{\geq 1} \setminus V_2$,
(viii) $\Delta = \cup_{n=1}^{\infty} (\Delta_{DC}(n) \cup \Delta_{DB}(n) \cup \Delta_{DA}(n))$.

Proof. (i) This is Theorem 1 **(e1)** for $k = 2$.

(ii) Theorem 1 **(e2)** for $k = 2$.

(iii) $\Delta_{DA}(n) = (h^3(n) - h^2(n)) + (h^2(n) - h(n)) = a^2(n) + a(n) = u^2(n)$ by Theorem 1 **(e1)** and **(b)**.

(iv) Theorem 1 **(c)**.

(v) Theorem 1 **(e1)**.

(vi) $\Delta_{BA}(n) : (a^3(n) - h^2(n)) + (h^2(n) - h(n)) = a(n) - 2$.

(vii) Notice that for every $n \in \mathbb{Z}_{\geq 1}$, $a^2(n) \in G_1$, $a^2(n) + 2 \in G_1$ (by Theorem 1 **(h1)**), $2 \in U_2$, $a(n) - 2 \in G_1 \cup U_2$ (by **(h3)**). It then follows from (iii) and (v) that $\Delta = G_1 \cup U_2$. The result follows since the sets G_1, U_2, V_2 clearly partition $\mathbb{Z}_{\geq 1}$.

(viii) Follows from (i)-(iii), (vii) and Theorem 1 **(h2)**. ■

Lemma 5. For fixed $n \in \mathbb{Z}_{\geq 1}$, let $0 < t < a^2(n)$, $t \notin V_2$. Then there exists $0 \leq m < n$ such that $t \in \Delta_{DC}(m) \cup \Delta_{DB}(m) \cup \Delta_{DA}(m)$.

Proof. We have $t < a^2(n) = \Delta_{DC}(n) < \Delta_{DB}(n) < \Delta_{DA}(n)$. It then follows from Lemma 4(viii), that there must be some $m < n$ for which $t \in \Delta_{DC}(m) \cup \Delta_{DB}(m) \cup \Delta_{DA}(m)$. ■

Theorem 2. The set T constitutes the set of P -positions of the game Flora.

Proof. To begin with we note the following facts:

- Lemma 1 implies that each of the sequences A_n, B_n, C_n, D_n is increasing (Lemma 1).
- A, B, C, D partition $\mathbb{Z}_{\geq 1}$ (Theorem 1 **(a)**).

It evidently suffices to prove the following two statements:

- (A) Every move from any position in T results in a position outside T .
- (B) For every position outside T there is a move into a position in T .

(A) Clearly there is no legal move $T_1 \rightarrow T_0$. Suppose that there are positions T_n, T_m with $m < n$, $n \geq 2$ such that there is a legal move $T_n \rightarrow T_m$. This move must be of type III, since A, B, C, D partition $\mathbb{Z}_{\geq 1}$, from which it follows easily, using Lemma 4, that $A_n < B_n < C_n < D_n$ for $n \geq 2$.

By Lemma 4(vii), $\Delta_{DC}(n) \notin V_2$, so we have to consider only move III(i). We first show that $D_n - p$ can only be D_m . It cannot be A_m , since then $C_n - p < A_m$ has no place in row m of T . Suppose $D_n - p = B_m$. Then $C_n - p = A_m$. But $\Delta_{BA}(m) < \Delta_{BA}(n) = a(n) - 2 < \Delta_{DC}(n) = a^2(n)$, contradicting the move rule III(i). Suppose $D_n - p = C_m$. Since $C_n - B_n = 2$ for all $n \in \mathbb{Z}_{\geq 1}$, we have $C_n - p = A_m$. But $\Delta_{CA}(m) = \Delta_{BA}(m) + 2 \leq \Delta_{BA}(n) + 1 = a(n) - 1 < \Delta_{DC}(n) = a^2(n)$, again contradicting move rule III(i). Thus indeed $D_n - p = D_m$.

Suppose $C_n - p = C_m$. Subtracting, $\Delta_{DC}(n) = \Delta_{DC}(m)$, so $a^2(n) = a^2(m)$ which is impossible for $m < n$ since the sequence $g^2(\ell)$ is strictly increasing.

Suppose $C_n - p = B_m = C_m - 2$. Then $\Delta_{DC}(n) = \Delta_{DB}(m) = \Delta_{DC}(m) + 2$ which is possible if and only if $p = 5$ and $m = n - 1$. But this case is excluded by the proviso. Finally, suppose that $C_n - p = A_m$. Then $\Delta_{DC}(n) = \Delta_{DA}(m)$. By Lemma 4 this is equivalent to $a^2(n) = u^2(m)$. This is possible for no $m < n$ by disjointness.

(B) Let $(a_1, a_2, a_3, a_4) \notin T$, $0 \leq a_1 \leq a_2 \leq a_3 \leq a_4$. If there is equality in any of these or $a_3 - a_2 = 1$, a move of type I or II leads to T_0 . So we may assume $0 < a_1 < a_2 < a_2 + 1 < a_3 < a_4$. By the complementarity of A, B, C, D , a_1 appears in precisely one component of precisely one T_n , $n \geq 1$. If $a_1 = D_n$, move $a_2 \rightarrow A_n, a_3 \rightarrow B_n, a_4 \rightarrow C_n$.

So suppose that $a_1 = C_n$.

If $a_4 \geq D_n$, move $a_2 \rightarrow A_n, a_3 \rightarrow B_n, a_4 \rightarrow D_n$. So assume $a_4 < D_n$. Let

$$t := a_4 - a_3.$$

We consider 2 cases.

(a) $t \notin V_2$, and (b) $t \in V_2$.

(a) $t \notin V_2$. We have

$$0 < t = a_4 - a_3 < D_n - a_3 < D_n - a_1 = D_n - C_n = \Delta_{DC}(n) = a^2(n).$$

By Lemma 5, there exists $m < n$ such that either (i) $t = \Delta_{DC}(m)$, or (ii) $t = \Delta_{DB}(m)$, or (iii) $t = \Delta_{DA}(m)$.

For case (i), move $(a_1, a_2, a_3, a_4) \rightarrow (A_m, B_m, C_m, D_m)$. This is a legal move: $a_1 = C_n > A_n > A_m$, and $a_2 > a_1 = C_n > B_n > B_m$, $a_3 > a_1 = C_n > C_m$, $a_4 = a_3 + D_m - C_m > D_m$, so this move (as well as in the remainder of this proof) is of the form III.

For case (ii), move $a_1 \rightarrow A_m, a_2 \rightarrow C_m, a_3 \rightarrow B_m, a_4 \rightarrow D_m$. This is a legal move: $a_1 = C_n > A_n > A_m$, $a_2 > a_1 = C_n > C_m$, $a_3 > a_1 = C_n > C_m > B_m$, $a_4 = a_3 + D_m - B_m > D_m$.

For case (iii), move $a_1 \rightarrow B_m, a_2 \rightarrow C_m, a_3 \rightarrow A_m, a_4 \rightarrow D_m$. This is a legal move: $a_1 = C_n > B_n > B_m$, $a_2 > a_1 = C_n > C_m$, $a_3 > a_1 = C_n > C_m > A_m$, $a_4 = a_3 + D_m - A_m > D_m$.

(b) $t \in V_2$. To remind ourselves, $t = a_4 - a_3$ and we have $a_1 = C_n, a_4 < D_n$. Now $t - 1 \in (V_2 - 1)$. Since $(V_2 - 1) \subset G_2$ (Theorem 1 **(h4)**), we have $t - 1 = a^2(m)$ for suitable $m \in \mathbb{Z}_{\geq 1}$. Also $\Delta_{DC}(m) = a^2(m)$ (Lemma 4 (i)). So we move: $(a_1, a_2, a_3, a_4) \rightarrow (A_m, B_m, C_m, D_m)$. This is a legal move:

- $m < n$, since $\Delta_{DC}(m) = a_4 - a_3 - 1 < D_n - a_1 = D_n - C_n = \Delta_{DC}(n)$.
- $a_1 = C_n > A_n > A_m, a_2 > a_1 = C_n > B_n > B_m, a_3 > a_1 = C_n > C_m, a_4 = a_3 + 1 + D_m - C_m > a_1 + D_m - C_m = C_n + D_m - C_m > D_m$.

So suppose that $a_1 = B_n$.

If $a_4 \geq D_n$, then move $a_2 \rightarrow A_n$, $a_3 \rightarrow C_n$, $a_4 \rightarrow D_n$. This is a legal move, since $a_2 > a_1 = B_n > A_n$ and

$$a_3 \geq a_2 + 1 \geq a_1 + 2 = B_n + 2 = C_n.$$

Therefore we may assume $a_4 < D_n$. The proof is similar to the above case $a_1 = C_n$. We have $a_3 \geq C_n$, and $0 < t - 1 < t = a_4 - a_3 < D_n - a_3 \leq \Delta_{DC}(n)$. Hence by Lemma 5 there is $m < n$ such that, for case (a), either (i) $t = \Delta_{DC}(m)$, or (ii) $t = \Delta_{DB}(m)$, or (iii) $t = \Delta_{DA}(m)$. For case (b) we have $t - 1 = a^2(m)$ for some $m \in \mathbb{Z}_{\geq 1}$.

(a) $t \notin V_2$.

For case (i), move $a_2 \rightarrow A_m$, $(a_3, a_4) \rightarrow (C_m, D_m)$. This is a legal move: $a_2 > a_1 = B_n > A_n > A_m$, $a_3 \geq C_n > C_m$,

$$a_4 = a_3 + D_m - C_m \geq C_n + D_m - C_m > D_m.$$

For case (ii), move $a_1 \rightarrow A_m$, $a_2 \rightarrow C_m$, $a_3 \rightarrow B_m$, $a_4 \rightarrow D_m$. This is a legal move: $a_1 = B_n > A_n > A_m$, $a_2 \geq a_1 + 1 = B_n + 1 = C_n - 1 \geq C_m$, $a_3 > a_1 = B_n > B_m$, $a_4 = a_3 + D_m - B_m \geq C_n + D_m - B_m > D_m$.

For case (iii), move $a_1 \rightarrow B_m$, $a_2 \rightarrow C_m$, $a_3 \rightarrow A_m$, $a_4 \rightarrow D_m$. This is a legal move: $a_1 = B_n > B_m$, $a_2 \geq a_1 + 1 = B_n + 1 = C_n - 1 \geq C_m$, $a_3 > a_1 = B_n > B_m > A_m$, $a_4 = a_3 + D_m - A_m > D_m$.

(b) $t \in V_2$. We have $t = a_4 - a_3$, $a_1 = B_n$, $a_4 < D_n$. As in case (b) above, we move $(a_1, a_2, a_3, a_4) \rightarrow (A_m, B_m, C_m, D_m)$. This is a legal move:

- $m < n$, since $\Delta_{DC}(m) = a_4 - a_3 - 1 < D_n - a_3 - 1 \leq D_n - C_n - 1 < \Delta_{DC}(n)$.
- $a_1 = B_n > A_n > A_m$, $a_2 > a_1 = B_n > B_m$, $a_3 \geq C_n > C_m$, $a_4 = a_3 + 1 + D_m - C_m \geq C_n + 1 + D_m - C_m > D_m$.

Finally, we consider the case $a_1 = A_n = h(n)$.

If $a_2 < h^2(n) - 2$ we can move to T_0 (rule (II)(iii)). Otherwise, $a_2 \geq h^2(n) - 2 = B_n$. Since $a_3 - a_2 > 1$, we have $a_3 \geq B_n + 2 = C_n$. If $a_4 \geq D_n$, then at least one of the inequalities for a_2, a_3, a_4 is strict, since $(a_1, a_2, a_3, a_4) \notin T$. Then move $(a_1, a_2, a_3, a_4) \rightarrow (A_n, B_n, C_n, D_n)$. If $a_4 < D_n$ then for case (a) there is $m < n$ such that $0 < t = a_4 - a_3 < \Delta_{DC}(n)$. Hence by Lemma 5, there is $m < n$ such that either (i) $t = \Delta_{DC}(m)$, or (ii) $t = \Delta_{DB}(m)$, or (iii) $t = \Delta_{DA}(m)$. For case (b), $0 < t - 1 = a^2(m)$ for some $m \in \mathbb{Z}_{\geq 1}$.

(a) $t \notin V_2$. For case (i) move $(a_1, a_2, a_3, a_4) \rightarrow (A_m, B_m, C_m, D_m)$. This is a legal move, since $m < n$ implies $a_1 = A_n > A_m$, $a_2 \geq B_n > B_m$, $a_3 \geq C_n > C_m$, $D_n > D_m$. For case (ii), move $a_1 \rightarrow A_m$, $a_2 \rightarrow C_m$, $a_3 \rightarrow B_m$, $a_4 \rightarrow D_m$. This is a legal move: $a_1 = A_n > A_m$, $a_2 \geq B_n = C_n - 2 > C_m$, where the strict inequality follows since $C_n - C_m \geq 3$ (Theorem 1 (f4) for $k = 3$). Also $a_3 \geq C_n > B_n > B_m$, $a_4 = a_3 + D_m - B_m > D_m$. For case (iii) move $a_1 \rightarrow B_m$,

$a_2 \rightarrow C_m, a_3 \rightarrow A_m, a_4 \rightarrow D_m$. We have to prove the legality of this move. We begin by showing that $a_1 = A_n > B_m$. Notice that

$$\begin{aligned} t &= a_4 - a_3 = \Delta_{DA}(m) = u^2(m) \quad (\text{Lemma 4}) \\ &< \Delta_{DC}(n) = a^2(n) \quad (\text{Lemma 4}) \\ &= h(n) - 1 \quad (\text{Theorem 1 (c)}). \end{aligned}$$

Thus $h(n) > u^2(m) + 1$. But $h(n) = A_n$ and $u^2(m) = a^3(m) + 1$ (by Theorem 1 (d)) $= B_m + 1$, so indeed $A_n > B_m + 2 > B_m$. Next,

$$\begin{aligned} a_2 &\geq B_n = a^3(n) = h^2(n) - 2 \quad (\text{Theorem 1 (d)}) \\ &> h^2(n - 1) \quad (\text{Theorem 1 (f4)}) = C_{n-1} \geq C_m. \end{aligned}$$

Also $a_3 \geq C_n > C_m > A_m$ and $a_4 = a_3 + D_m - A_m \geq C_n + D_m - A_m > D_m$.

(b) $t \in V_2$. We have $t = a_4 - a_3, a_1 = A_n, a_4 < D_n, t - 1 = a^2(m) = \Delta_{DC}(m)$ for some $m \in \mathbb{Z}_{\geq 1}$. As above we move $(a_1, a_2, a_3, a_4) \rightarrow (A_m, B_m, C_m, D_m)$. This is a legal move:

- $m < n$, since $\Delta_{DC}(m) = a_4 - a_3 - 1 < D_n - C_n - 1 < \Delta_{DC}(n)$.
- $a_1 = A_n > A_m, a_2 \geq B_n > B_m, a_3 \geq C_n > C_m, a_4 = a_3 + 1 + D_m - C_m > C_n + D_m - C_m > D_m$. ■

Theorem 3. *The algebraic winning strategy of Flora precipitates a polynomial-time algorithm for consummating a win.*

Proof. Given a position (a_1, a_2, a_3, a_4) of Flora with $0 \leq a_1 \leq a_2 \leq a_3 \leq a_4$. Its input size is $O(\log a_1 + \log a_2 + \log a_3 + \log a_4)$. Whether or not move rules II(i) or II(ii) apply can be checked trivially. We know (Theorem 1 (a)) that a_1 is precisely one of $a^3(n), h^3(n), h^2(n), h(n)$. We have to find out which it is, and the corresponding value of n .

Suppose first that $a_1 = a^3(n) = \lfloor \varphi \lfloor \varphi \lfloor n \varphi \rfloor \rfloor \rfloor$. Using the inequality $x - 1 < \lfloor x \rfloor \leq x$, a straightforward computation shows that $\lfloor a_1 \varphi^{-3} \rfloor + 1 \leq n \leq \lfloor (a_1 + 1) \varphi^{-3} \rfloor + 1$. Computing φ to $O(\log a_1)$ places gives the range for the candidate values of n and for each of them (one or two), we have to compute $a^3(n)$, comparing it with a_1 . A similar computation can be done for $h^3(n), h^2(n), h(n)$. (Notice that there is not necessarily an integer candidate n for some of these ranges. For example, if we suppose that $a_1 = h^3(n)$, then we get $\lfloor a_1 \varphi^{-4} \rfloor + 1 \leq n \leq \lfloor a_1 \varphi^{-4} + 2 \varphi^{-2} \rfloor$.) The same method also indicates whether or not $a_4 - a_3 \in V_2$, or whether move rule II(iii) applies. All these computations can be done in linear time in the input size.

Finally, we use a binary search to find $m \in [0, n]$ such that if $a_4 - a_3 \notin V_2$, then $h^3(m) - h^2(m) = a_4 - a_3$ or some other difference of the columns in the m -th row is $a_4 - a_3$. Similarly for the case $a_4 - a_3 \in V_2$. ■

3.2 Recursive Formulation of the P -positions

Let $S \subsetneq \mathbb{Z}_{\geq 1}$ and $\bar{S} = \mathbb{Z}_{\geq 1} \setminus S$. The “Minimum EXcludant” of S is defined by

$$\text{mex } S = \min \bar{S} = \text{least positive integer not in } S.$$

In particular, the mex of the empty set is 1. (This somewhat nonstandard definition of the mex function is needed for §5.)

Let $T'_0 = (0, 0, 0, 0)$, $T'_1 = (2, 1, 3, 4)$. If $T'_m := (A'_m, B'_m, C'_m, D'_m)$ has already been defined for all $m < n$ ($n \geq 2$), then let

$$A'_n = \text{mex}\{A'_i, B'_i, C'_i, D'_i : 0 \leq i < n\},$$

$$B'_n = \begin{cases} B'_{n-1} + 3 & \text{if } A'_n - A'_{n-1} = 2 \\ B'_{n-1} + 5 & \text{otherwise,} \end{cases}$$

$$C'_n = B'_n + 2,$$

$$D'_n = \begin{cases} D'_{n-1} + 5 & \text{if } A'_n - A'_{n-1} = 2 \\ D'_{n-1} + 8 & \text{otherwise.} \end{cases}$$

Let $T' := \cup_{n=0}^{\infty} T'_n$.

Theorem 4. *The set T' constitutes the set of P -positions of the game Flora.*

Proof. We show that for all $m \in \mathbb{Z}_{\geq 0}$, $A'_m = A_m$, $B'_m = B_m$, $C'_m = C_m$, $D'_m = D_m$. Suppose this holds for all $m < n$ ($n \geq 1$). Let $E = \text{mex}\{A_i, B_i, C_i, D_i : 0 \leq i < n\}$. The value E cannot have been assumed in any of the four sequences for $m < n$, since A, B, C, D split $\mathbb{Z}_{\geq 1}$, so $E \geq A_n$. If $E > A_n$, then A_n would never be assumed since the sequences are strictly increasing, again contradicting the complementarity of the sequences. Thus $A_n = E = A'_n$, and the other 3 equalities follow from Theorem 2 (f). ■

The definition of the set T' is straightforward; it doesn't use the functions $h(n)$, $a(n)$ used for defining T . Thus the recursive computation of T' looks easier than that of the set T . Moreover, the proof of Theorem 4 is very short, and that of Theorem 2 is long.

However, the proof of Theorem 4 leans heavily on Theorems 2 and 1. If the initial position of the game is (a_1, a_2, a_3, a_4) , the input size is $\log a_1 + \log a_2 + \log a_3 + \log a_4$. The time needed to compute whether the position is a P -position or not, seems to be proportional to $a_1 + a_2 + a_3 + a_4$, because the unwieldy mex function appears to require scanning previous entries of the sequences A_n, B_n, C_n, D_n . However, a new method [12] shows that actually also the algorithm implied by Theorem 4 is polynomial.

3.3 Arithmetic Formulation of the P -positions

For $N \in \mathbb{Z}_{\geq 1}$, let $R(N) = (d_m, \dots, d_0)$ be the representation of N in the Fibonacci numeration system (recall Notation 2). Then $(d_m, \dots, d_0, 0)$ is the *left shift* of $R(N)$.

Theorem 5. $R(A)$ is the set of all representations that end in an odd number of 0-bits in the Fibonacci numeration system, $R(B)$ the set of all representations that end in 001, $R(C)$ the set of all representations that end in a positive even number of 0-bits, and $R(D)$ the set of all representations that end in 101. Moreover, for every $n \in \mathbb{Z}_{\geq 1}$, $R(C_n)$ is the left shift of $R(A_n)$.

See Table 3.3 for an example.

Table 3.3: Representation of the P -positions in the Fibonacci numeration system.

21	13	8	5	3	2	1	A_n	n	B_n	34	21	13	8	5	3	2	1
					1	0	2	1	1								1
			1	0	0	0	5	2	6					1	0	0	1
			1	0	1	0	7	3	9				1	0	0	0	1
		1	0	0	1	0	10	4	14			1	0	0	0	0	1
	1	0	0	0	0	0	13	5	19			1	0	1	0	0	1
	1	0	0	0	1	0	15	6	22		1	0	0	0	0	0	1
	1	0	1	0	0	0	18	7	27		1	0	0	1	0	0	1
	1	0	1	0	1	0	20	8	30		1	0	1	0	0	0	1
1	0	0	0	0	1	0	23	9	35	1	0	0	0	0	0	0	1
1	0	0	1	0	0	0	26	10	40	1	0	0	0	1	0	0	1

34	21	13	8	5	3	2	1	C_n	n	D_n	55	34	21	13	8	5	3	2	1	
					1	0	0	3	1	4								1	0	1
			1	0	0	0	0	8	2	12					1	0	1	0	1	
			1	0	1	0	0	11	3	17				1	0	0	1	0	1	
		1	0	0	1	0	0	16	4	25			1	0	0	0	1	0	1	
	1	0	0	0	0	0	0	21	5	33			1	0	1	0	1	0	1	
	1	0	0	0	1	0	0	24	6	38		1	0	0	0	0	1	0	1	
	1	0	1	0	0	0	0	29	7	46		1	0	0	1	0	1	0	1	
	1	0	1	0	1	0	0	32	8	51		1	0	1	0	0	1	0	1	
1	0	0	0	0	1	0	0	37	9	59	1	0	0	0	0	0	1	0	1	
1	0	0	1	0	0	0	0	42	10	67	1	0	0	0	1	0	1	0	1	

Proof. The proof is similar to that of Theorem 3 of [8]. For every $m \in \mathbb{Z}_{\geq 1}$, $R(\lfloor m\varphi \rfloor)$ ends in an even number of 0-bits (including 0 0-bits), and $R(\lfloor m\varphi^2 \rfloor)$

ends in an odd number of 0-bits ([5], §4). Hence $R(A)$ is the set of all numbers that end in an odd number of 0-bits in the Fibonacci numeration system, whereas each of the other 3 representations ends in an even number of 0-bits. Now $R(C)$ is the set of all numbers that end in a positive even number of 0-bits ([8]), hence $R(B)$ and $R(D)$ each end in a 1-bit. Recall that $C_n = B_n + 2$. If $R(B)$ would contain a number with representation ending in 101, then adding 2 to it would end in 1 (since $2+3=5$ is the next Fibonacci number), contradicting the form of $R(C_n)$. Therefore $R(B)$ is the set of all numbers ending in 001. By complementarity, $R(D)$ is therefore the set of all numbers ending in 101.

Since $R(A)$ is the set of all representations ending in an odd number of 0-bits, and $R(C)$ is the set of all representations ending in a positive even number of 0-bits, the latter is the left shift of the former. Suppose that $R(C_m)$ is the left shift of $R(A_m)$ for every $m < n$. If $R(C_n)$ would not be the left shift of $R(A_n)$, then it would be assumed later on (by complementarity), contradicting the strict increase of C . ■

This formulation of the P -positions is also easily seen to lead to a polynomial-time winning strategy.

4 The Word-mapping Approach

In this section we show how to construct G_k recursively by a word-map for every $k \in \mathbb{Z}_{\geq 1}$. Similar methods can be used to construct other functions defined in §2. We also present our fourth, word-mapping, formulation of the P -positions of the game Flora. The length of any (sub)word w is denoted by $|w|$.

4.1 Word-mapping for G_k

Define the morphism $1 \rightarrow 10, 0 \rightarrow 1$. Its fixed point is the word: $F = 1011010110110\dots$, also known as the Fibonacci word. For $k \geq 1$, the characteristic function χ_k of G_k is defined by

$$\chi_k(m) = \begin{cases} 1 & \text{if } \exists n \text{ s.t. } a^k(n) = m \\ 0 & \text{otherwise.} \end{cases}$$

Definition 1. Given a binary word W . A *run* of 0s is any (possibly empty) subword of W consisting solely of 0s, flanked on the left and right by a 1-bit. A *block* is any subword consisting of a 1-bit followed by a run of 0s.

Theorem 6. For every $k \geq 1$, the word-mapping for producing the characteristic function χ_k of G_k , beginning with $10^{F_{k-1}-1}$, is:

$$10^{F_{k-1}-1} \rightarrow 10^{F_k-1}, \quad 0^{F_{k-2}} \rightarrow 10^{F_{k-1}-1}.$$

Proof. Notice that for G_1 , the word-mapping is simply the well-known morphism $1 \rightarrow 10, 0 \rightarrow 1$, which produces F . Moreover, $\chi_1 = F$. See e.g., [1], ch. 9 and [11].

The word-map is well-defined. Indeed, the initial block of length F_{k-1} is mapped into a block B_1 of length F_k . In the second iteration, the prefix of length F_{k-1} of B_1 is again mapped into B_1 . The remaining abutting suffix of B_1 consists of F_{k-2} 0s, so it is mapped into a block B_2 of length F_{k-1} . In the third iteration, B_1 and B_2 are generated again, and then the block B_2 of length F_{k-1} generates a block B_1 . Thus for all subsequent iterations only blocks of the form B_1 and B_2 are generated, and there is never any parsing conflict.

Since $\chi_1 = 1011010110110\dots$, Theorem 1(f) implies that $\chi_2 = 100101001001010010100\dots$, where we inserted into χ_1 $F_1 - F_0 = 1$ zero to each run of $F_0 - 1 = 0$ zeros (i.e., one 0 between every consecutive 1s), and $F_2 - F_1 = 1$ zero to each run of $F_1 - 1 = 1$ zeros. Doing this yields distances between consecutive 1s in χ_2 of F_1 and F_2 , precisely at the locations where the distances between consecutive 1s of χ_1 are F_0 and F_1 respectively. Similarly, $\chi_3 = 1000010010000100001001000010010000\dots$, where we inserted into χ_2 $F_2 - F_1 = 1$ zero to each run of $F_1 - 1 = 1$ zero, and $F_3 - F_2 = 2$ zeros to each run of $F_2 - 1 = 2$ zeros.

In general, for producing χ_{k+1} from χ_k , we add to χ_k $F_k - F_{k-1} = F_{k-2}$ zeros to each run of $F_{k-1} - 1$ zeros and $F_{k+1} - F_k = F_{k-1}$ zeros to each run of F_k zeros. This yields blocks of sizes F_k and F_{k+1} respectively, at the locations specified by Theorem 1(f).

Assume inductively that the word-mapping

$$10^{F_{k-1}-1} \rightarrow 10^{F_k-1}, \quad 0^{F_{k-2}} \rightarrow 10^{F_{k-1}-1}$$

produces χ_k , so it generates distances between consecutive 1s of F_{k-1} and F_k at the locations specified by Theorem 1(f). Then the word-mapping

$$10^{F_k-1} \rightarrow 10^{F_{k+1}-1}, \quad 0^{F_{k-1}} \rightarrow 10^{F_k-1}$$

produces χ_{k+1} , since it adds $F_{k+1} - F_k$ to the F_k 0s of the long 0-runs of χ_k , and $F_k - F_{k-1}$ 0s to the F_{k-1} short 0-runs of χ_k . ■

4.2 Word-mapping Formulation of of the P -positions

Denote terms of A_n, B_n, C_n, D_n by a, b, c, d respectively.

Theorem 7. *The word-mapping*

$$bac \rightarrow bacda, \quad da \rightarrow bac,$$

beginning with bac , generates the characteristic function of the P -positions of the Flora game.

Proof. The proof is rather similar to that of Theorem 6, and is therefore omitted. ■

This theorem also leads to a polynomial-time winning strategy, since induction shows that for every $k \in \mathbb{Z}_{\geq 1}$, the k -th application of the word-mapping generates a word of length F_{k+2} .

Notice that if we replace bac by 1 and da by 0, we get back our old morphic friend $1 \rightarrow 10, 0 \rightarrow 1$.

4.3 Characterization of the Sequences G , H , by the Fibonacci Numeration System

We know from Lemma II and §3.3, that $R(a(n))$ ends in an even number of 0s, $R(h(n))$ in an odd number of 0s, $R(h^2(n))$ in an even positive number of 0s, $R(h^3(n))$ in 101, and $R(g^3(n))$ in 10^s1 , $s \geq 2$. What's the general pattern?

Theorem 8. (i) $R(G_1)$ is the set of all representations that end in an even number of 0s, $R(H_1)$ is the set of all representations that end in an odd number of 0s, $R(G_2)$ is the set of all representations that end in a 1-bit, and $R(H_2)$ is the set of all representations that end in an even positive number of 0s.

(ii) For every $n \in \mathbb{Z}_{\geq 1}$, $R(h(n))$ is the left shift of $R(a(n))$; and $R(h^2(n))$ is the left shift of $R(h(n))$.

(iii) For every $k \in \mathbb{Z}_{\geq 3}$ and all $n \in \mathbb{Z}_{\geq 1}$, $R(G_k)$ is the set of all representations that end in the word 10^s1 for all $s \geq k-1$, and $R(H_k)$ is the set of all representations that end in the word $10^{k-2}1$ (left 1-bit in position $k-1$).

Proof. Items (i) and (ii) are already known from Theorem 5 and Lemma II, and are included here only for the sake of completeness. We only have to point out the statement about $R(G_2)$, which follows from the fact that G_2, H_2, H_1 split the positive integers (see also [8], Theorem 3).

(iii) Induction on k . The base case $k=3$ was proved in Theorem 5. For $k \geq 3$, suppose that we already proved that $R(G_k)$ is the set of all representations that end in 10^s1 for all $s \geq k-1$, and $R(H_k)$ is the set of all representations that end in $10^{k-2}1$. It clearly remains only to show that $R(G_{k+1})$ is the set of all representations that end in 10^s1 for all $s \geq k$, and $R(H_{k+1})$ is the set of all representations that end in the word $10^{k-1}1$. Recall Theorem 1 (c): $h^k(n) = g^{k+1}(n) + F_{k-1}$. If $R(G_{k+1})$ would contain a number, say $a^{k+1}(n)$, with representation ending in $10^{k-1}1$ (with leftmost 1-bit in position k), then adding F_{k-1} to it would result in a word with representation ending in 0^k1 because $F_{k-1} + F_k = F_{k+1}$ is the next Fibonacci number. But then $R(h^k(n))$ would end in 10^s1 for some $s \geq k$, contradicting the induction hypothesis. Thus $R(H_{k+1})$ is the set of all representations that end in the word $10^{k-1}1$. Since $G_{k+1}, H_{k+1}, H_k, \dots, H_2, H_1$ split the integers, $R(H_{k+1})$ is the set of all representations that end in the word $10^{k-1}1$. ■

Notes. (1) The general pattern of the representation of the suffixes of H_k for $k \geq 3$ is quite different from that of H_1 and H_2 , and both of these are different from each other. The same holds for G_k , $k \geq 3$ and G_1 and G_2 . Therefore the induction proof could not have begun with $k=1$ or 2.

(2) The statement in (i) about $R(G_1)$ and $R(H_1)$ is Theorem 9.1.15 (see also Corollary 9.1.14) in [1], credited there to [10]. (It is also Lemma II above.) The proof method of [1] follows [3].

Membership Problem. Fix some $k \in \mathbb{Z}_{\geq 1}$. Given $N \in \mathbb{Z}_{\geq 1}$. Then N is in precisely one of $G_k, H_k, H_{k-1}, \dots, H_2, H_1$ (Theorem 1 (a)). Can the following problem be solved in polynomial time?

Problem. Determine the set in which N lies.

Corollary 1. For every $k \in \mathbb{Z}_{\geq 1}$, the membership problem can be solved in linear time.

Proof. This can be proved by generalizing the method for computing n in the proof of Theorem 3 to the case of arbitrary k . But a more “elegant” method is to compute the Fibonacci representation of N , which can be done in linear time in the input size $\Theta(\log N)$. Theorem 8 then implies that the membership problem can be solved by scanning the suffix of $R(N)$, at most all of its $\Theta(\log N)$ bits. ■

5 Infinite Complementary Arrays

The doubly infinite *Stolarsky Array* A with entries $A(i, j)$, $i, j \geq 1$ [22], is defined as follows: For every $m \geq 1$, $A(m, 1) = \text{mex}\{A(i, j) : i < m, j \geq 1\}$, $A(m, 2) = \lfloor \varphi A(m, 1) + 1/2 \rfloor$, and for all $i \geq 1$, $j \geq 3$, $A(i, j) = A(i, j - 1) + A(i, j - 2)$. Then every positive integer appears precisely once in A . A beginning portion is exhibited in Table 2. Many variations, interspersions and dispersions have since been given, see e.g., [16], [19]. All are doubly infinite, $\lim_{j \rightarrow \infty} (A(i, j + 1) - A(i, j)) = \infty$ for every $i \geq 1$, and every positive integer appears precisely once in A .

Table 2: The Stolarsky array

1	2	3	5	8	13	...
4	6	10	16	26	42	...
7	11	18	29	47	76	...
9	15	24	39	63	102	...
⋮	⋮	⋮	⋮	⋮	⋮	⋮

For every $k \geq 1$, define the Flora-Array L_K with the $k+1$ rows $H_1, H_2, \dots, H_k, G_k$. This array has a different character. By Theorem 1 (a), this *singly*-infinite array also has the property that every positive integer appears precisely once. Moreover, $A(i, j + 1) - A(i, j) \in \{F_i, F_{i+1}\}$ is bounded for every fixed i and all $j \geq 1$, but $\lim_{i \rightarrow \infty} (A(i, j + 1) - A(i, j)) = \infty$. Table 3 depicts the case $k = 6$. The two bottom lines, below the horizontal line, illustrate the fact that G_7, H_7 split G_6 , so replacing G_6 by H_7 and G_7 constitutes L_7 .

Table 3: A complementary Flora array L_6 with 7 rows

n	1	2	3	4	5	6	7	8	9	10	11	12	13
H_1	2	5	7	10	13	15	18	20	23	26	28	31	34
H_2	3	8	11	16	21	24	29	32	37	42	45	50	55
H_3	4	12	17	25	33	38	46	51	59	67	72	80	88
H_4	6	19	27	40	53	61	74	82	95	108	116	129	142
H_5	9	30	43	64	85	98	119	132	153	174	187	208	229
H_6	14	48	69	103	137	158	192	213	247	281	302	336	370
G_6	1	22	35	56	77	90	111	124	145	166	179	200	221
H_7	22	77	111	166	221	255	310	344	...				
G_7	1	35	56	90	124	145	179	200	234	268	289	323	...

6 Epilogue

We have generated sequences consisting of nested arbitrary applications of the floor function to φ and φ^2 , established many identities and relationships involving them and then applied them to formulate an algebraic winning strategy to the game Flora. We also presented recursive, arithmetic and word-mapping formulations of the winning strategy. In addition, we characterized the main sequences by means of the Fibonacci numeration system, and generated infinite complementary arrays of the sequences.

Can some of the relationships of the iterated floor functions be generalized to reals other than φ and φ^2 ? As a first step, we could interchange φ with φ^2 , studying the ensuing sequences and the games implied by them. Specifically, define $g'(n) = \lfloor n\varphi^2 \rfloor$, $h'(n) = \lfloor n\varphi \rfloor$, and for $k \geq 2$, $g'^k(n) = \lfloor \varphi^2 g'^{k-1}(n) \rfloor$, $h'^k(n) = \lfloor \varphi^2 g'^{k-1}(n) \rfloor$. If we define G'_i, H'_i in the obvious way, it's straightforward to see that then the first item of Theorem 1, is preserved, namely $G'_k, H'_k, H'_{k-1}, \dots, H'_2, H'_1$ partition $\mathbb{Z}_{\geq 1}$. What games can be spawned from this partition?

More generally, it would be well to investigate which of the above results hold for which classes of positive reals beyond φ . For example, Lemma 1 and Theorem 1(a) clearly hold if we replace φ by any irrational $\alpha \in (1, 2)$ and φ^2 by $\beta = \alpha/(\alpha - 1)$. Perhaps large parts of Theorem 1 can be generalized for the case where $\alpha = (2 - t + \sqrt{t^2 + 4})/2$, $\beta = \alpha + t$, where t is any given positive integer, since then the simple continued fraction of α is $[1, t, t, t, \dots]$, so the numeration system arguments used in the proof of (c) and (d) of Theorem 1 carry over in a simple way. What games are induced by these relationships?

The notion of arbitrary iterations of the floor function appeared in [21] and [15]. In the former, the iterations are with *rational* numbers whose sizes depend on the iteration depth; in the latter, the aim is to represent the positive integers in the form of iterated floor functions involving φ and φ^2 . These are quite different from our iterations of the floor function. However, in [18] iterations of the form considered here were studied as pointed out above. See also [17].

The Raleigh game [8] is a 3-pile extension — not generalization — of Wythoff’s game. Flora is an extension of Raleigh. Although Flora appears not to be decomposable into sums of more elementary games, we were able to formulate for it three polynomial-time winning strategies. The one based on the Fibonacci numeration system is of particular interest. It demonstrates once again that numeration systems can make strategies of games in PrimGame efficient, similarly to appropriate data structures — see [20].

We can also define a 5-pile extension of Flora, but in the sequence of games with increasing number of piles, both the definition of the games and the validity proof of their strategies seem to become more difficult. For example, whereas the union of the differences Δ between the 3 columns of the P -positions of Raleigh covers all of $\mathbb{Z}_{\geq 1}$, the same union for the four columns of the P -positions of Flora leaves out V_2 . But perhaps a pattern for these games will emerge. This possibility may not be so far-fetched, since, as we saw, e.g., in §4.3, the general behavior begins only with $k = 3$ (corresponding to a game with 4 piles).

Acknowledgment. Thanks to Herb Wilf, who had written to me that he defined the sequence $a^k(n)$ for solving part (a) of AMM problem 11238 (113 Aug-Sept 2006, 655; Sol. 115 Aug-Sept 2008, 667), and conjectured that $\Delta a^k(n)$ assumes only two values for every fixed $k, n \in \mathbb{Z}_{\geq 1}$. His communication and earlier work of mine – including joint work with Eric Duchêne, Richard Nowakowski and Michel Rigo, to appear in JCT-A – motivated this paper.

References

- [1] J-P. Allouche and J. Shallit [2003], *Automatic Sequences: Theory, Applications, Generalizations*, Cambridge University Press, Cambridge.
- [2] E.R. Berlekamp, J.H. Conway and R.K. Guy [2001-2004], *Winning Ways for your Mathematical Plays*, Vol. 1-4, A K Peters, Wellesley, MA, 2nd edition: vol. 1 (2001), vols. 2, 3 (2003), vol. 4 (2004).
- [3] T.C. Brown [1993], Descriptions of the characteristic sequence of an irrational, *Canad. Math. Bull.* **36**, 15–21.
- [4] E. Duchêne and M. Rigo [2008], A morphic approach to combinatorial games: the Tribonacci case, *Theor. Inform. Appl.* **42**, 375–393.
- [5] A.S. Fraenkel [1982], How to beat your Wythoff games’ opponent on three fronts, *Amer. Math. Monthly* **89**, 353–361.
- [6] A.S. Fraenkel [1985], Systems of numeration, *Amer. Math. Monthly* **92**, 105–114.
- [7] A.S. Fraenkel [2004], Complexity, appeal and challenges of combinatorial games, *Theoret. Comput. Sci.* **313**, 393–415.

- [8] A.S. Fraenkel [2007], The Raleigh game, in: *Combinatorial Number Theory*, Proc. Integers Conference 2005, in celebration of the 70th birthday of Ronald Graham, Carrollton, Georgia, Oct. 27-30, 2005; B. Landman, M. Nathanson, J. Nešetřil, R. Nowakowski, C. Pomerance, eds., de Gruyter, pp. 199-208, 2007; appeared also in *Integers, Electr. J. of Combinat. Number Theory* **7(2)**, special volume in honor of Ron Graham, #A13, 11 pp., 2007 <http://www.integers-ejcnt.org/>
- [9] A.S. Fraenkel and D. Krieger [2004], The structure of complementary sets of integers: a 3-shift theorem, *Internat. J. Pure and Appl. Math.* **10**, 1–49.
- [10] A.S. Fraenkel, J. Levitt and M. Shimshoni [1972], Characterization of the set of values $f(n) = [n\alpha]$, $n = 1, 2, \dots$, *Discrete Math.* **2**, 335–345.
- [11] A.S. Fraenkel, M. Mushkin and U. Tassa [1978], Determination of $[n\theta]$ by its sequence of differences, *Canad. Math. Bull.* **21**, 441–446.
- [12] A.S. Fraenkel and U. Peled, Harnessing the unwieldy MEX function, Preprint.
- [13] G.H. Hardy [1999], *Ramanujan: Twelve Lectures on Subjects Suggested by his Life and Work*, AMS Chelsea Publishing, Providence, RI, Reprinted.
- [14] G.H. Hardy and E.M. Wright [1960], *An Introduction to the Theory of Numbers*, 4th edition, Oxford, UK.
- [15] V.E. Hoggatt and M. Bicknell-Johnson [1979], Representations of integers in terms of greatest integer functions and the golden section ratio, *Fibonacci Quart.* **17**, 306–318.
- [16] C. Kimberling [1993], Interspersions and dispersions, *Proc. Amer. Math. Soc.* **117**, 313–321.
- [17] C. Kimberling [2007], Complementary equations, *J. Integer Seq.* **10**, 14 pp. (electronic).
- [18] C. Kimberling [2008], Complementary equations and Wythoff sequences, *J. Integer Seq.* **11**, 5 pp. (electronic).
- [19] D.R. Morrison [1980], A Stolarsky array of Wythoff pairs, in: *A collection of manuscripts related to the Fibonacci sequence*, Fibonacci Assoc., Santa Clara, Calif., pp. 134–136.
- [20] B. Obrenić, Hyperbaric numeration systems and improved Fibonacci coding, Preprint.
- [21] K.A. Redish and W.F. Smyth [1991], Closed form expressions for the iterated floor function, *Discrete Math.* **91**, 317–321.
- [22] K.B. Stolarsky [1977], A set of generalized Fibonacci sequences such that each natural number belongs to exactly one, *Fibonacci Quart.* **15**, 224.

- [23] X. Sun [2005], Wythoff's sequence and N -heap Wythoff's conjectures, *Discrete Math.* **300**, 180–195.
- [24] X. Sun and D. Zeilberger [2004], On Fraenkel's N -heap Wythoff's conjectures, *Ann. Comb.* **8**, 225–238.