

(2005/12/17)

A quenched CLT for super-Brownian motion with random immigration

Wenming Hong¹ and Ofer Zeitouni²

Abstract. A quenched central limit theorem is derived for the super-Brownian motion with super-Brownian immigration, in dimension $d \geq 4$. At the critical dimension $d = 4$, the quenched and annealed fluctuations are of the same order but are not equal.

Key words: Super-Brownian motion, quenched central limit theorem, random immigration .

Mathematics Subject Classification (2000): Primary 60J80; Secondary 60F05.

1 Introduction and statement of results

Super-Brownian motion with super-Brownian immigration (SBMSBI, for short), is a superprocess in random environment, where the environment is determined by an immigration process which is controlled by the trajectory of another super-Brownian motion. Many interesting limit properties for SBMSBI were described under the *annealed* probability ([H02], [H03], [HL99] and [Zh05]). In this paper, we study the central limit theorem (CLT) under the *quenched* probability, that is, conditioned upon a realization of the immigration process, for $d \geq 4$.

To state our results and explain our motivation, we begin by recalling the SBMSBI model (we refer to [D93] and [P02] for a general introduction to the theory of superprocesses). Let $C(\mathbb{R}^d)$ denote the space of continuous bounded functions on \mathbb{R}^d . We fix a constant $p > d$ and let $\phi_p(x) := (1 + |x|^2)^{-p/2}$ for $x \in \mathbb{R}^d$. Let $C_p(\mathbb{R}^d) := \{f \in C(\mathbb{R}^d) : \sup |f(x)|/\phi_p(x) < \infty\}$. Let $M_p(\mathbb{R}^d)$ be the space of Radon measures μ on \mathbb{R}^d such that $\langle \mu, f \rangle := \int f(x)\mu(dx) < \infty$ for all $f \in C_p(\mathbb{R}^d)$. We endow $M_p(\mathbb{R}^d)$ with the p -vague topology, that is, $\mu_k \rightarrow \mu$ if and only if $\langle \mu_k, f \rangle \rightarrow \langle \mu, f \rangle$ for all $f \in C_p(\mathbb{R}^d)$. Then $M_p(\mathbb{R}^d)$ is metrizable ([I86]). We denote by λ the Lebesgue measure on \mathbb{R}^d , and note that $\lambda \in M_p(\mathbb{R}^d)$.

Let $S_{s,t}$ denote the heat semigroup in \mathbb{R}^d , that is, for $t > s$ and $f \in C(\mathbb{R}^d)$,

$$S_{s,t}f(x) = \frac{1}{(2\pi(t-s))^{d/2}} \int_{\mathbb{R}^d} e^{-|y-x|^2/2(t-s)} f(y) dy.$$

We write $S_t := S_{0,t}$ and G for the corresponding potential operator, that is $Gf = \int_0^\infty S_t f dt$, omitting the space variable x from the notation when no confusion may occur. Given $\mu \in$

¹School of Mathematical Science, Beijing Normal University, Beijing, 100875, P. R. China. Supported by NSFC (Grant No. 10121101) and the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry. Email: wmhong@bnu.edu.cn; wmhong@umn.edu.

²Department of Mathematics, University of Minnesota and Depts. of EE and Math., Technion. This work was partially supported by NSF grant DMS-0503775.

$M_p(\mathbb{R}^d)$, a *super-Brownian motion* $\varrho = (\varrho_t, P_\mu)$ is an $M_p(\mathbb{R}^d)$ -valued Markov process with $\varrho_0 = \mu$ and Laplace transform given by

$$E_\mu \exp\{-\langle \varrho_t, f \rangle\} = \exp\{-\langle \mu, v(t, \cdot) \rangle\}, \quad f \in C_p^+(\mathbb{R}^d), \quad (1.1)$$

where $v(\cdot, \cdot)$ is the unique mild solution of the evolution equation

$$\begin{cases} \dot{v}(t) = \frac{1}{2}\Delta v(t) - v^2(t) \\ v(0) = f, \end{cases} \quad (1.2)$$

and E_μ denotes expectation with respect to P_μ .

Given a super-Brownian motion $\varrho = (\varrho_t, P_\mu)$ as the “environment”, we will consider another super-Brownian motion with the immigration rate controlled by the trajectory of ϱ , the (SBMSBI) $X^\varrho = (X_t^\varrho, P_\nu^\varrho)$ with $X_0^\varrho = \nu$, which is again an $M_p(\mathbb{R}^d)$ -valued Markov process whose *quenched* probability law is determined by

$$E_\nu^\varrho \exp\{-\langle X_t^\varrho, f \rangle\} = \exp\{-\langle \nu, v(t, \cdot) \rangle - \int_0^t \langle \varrho_s, v(t-s, \cdot) \rangle ds\}. \quad (1.3)$$

Again, E_ν^ϱ denotes expectations with respect to P_ν^ϱ .

In the following we take $\mu = \nu = \lambda$, and write P^ϱ (resp. P) for P_λ^ϱ (resp. P_λ). We also use E^ϱ and E for the corresponding expectations. This model was considered in [HL99] and [H02, H03], see also [DGL02], where some interesting and new phenomena were revealed under the *annealed* probability law:

$$\mathbb{P}(\cdot) := \int P^\varrho(\cdot) P(d\varrho)$$

with expectation denoted by \mathbb{E} .

Our motivation for the present study is the annealed CLT derived in [HL99], which is summarized in Theorem 1.1 below.

Theorem 1.1 (Hong-Li) *Set*

$$\bar{a}_d(T) = \begin{cases} T^{3/4}, & d = 3 \\ T^{1/2}, & d \geq 4, \end{cases}$$

and with $f \in C_p^+(\mathbb{R}^d)$, define

$$\bar{Z}_T^\varrho(f) := \bar{a}_d(T)^{-1} \{\langle X_T^\varrho, f \rangle - \mathbb{E}\langle X_T^\varrho, f \rangle\},$$

Then, $\bar{Z}_T^\varrho(f) \Rightarrow \bar{Z}_\infty(f)$ in distribution under the law \mathbb{P} as $T \rightarrow \infty$, where $\bar{Z}_\infty(f)$ is a zero mean Gaussian random variable of variance

$$\text{var}(\bar{Z}_\infty(f)) = \begin{cases} \langle \lambda, f \rangle^2 / 6\pi^{3/2}, & d = 3, \\ \langle \lambda, f \rangle^2 / 8\pi^2 + \langle \lambda, fGf \rangle / 2, & d = 4, \\ \langle \lambda, fGf \rangle / 2, & d \geq 5. \end{cases}$$

In particular, contrasting with the standard super Brownian motion ([I86, Theorem 5.5 and Remark 6.1]), the SBMSBI exhibits smoothing of the critical dimension $d = 4$, since a logarithmic term is missing in the description of the long time behavior.

In the study of motion in random media, differences exist between quenched and annealed CLT behavior, and this difference is often tied to dimension and vanishes for dimension above

some critical value. See [Ze04] and [RS05] for several examples. It is thus of interest to identify whether a similar behavior occurs in the case of SBMSBI. Our main result, Theorem 1.2 below, shows that this is indeed the case.

Define the centered functional $Z_T^{\varrho}(f)$ by

$$Z_T^{\varrho}(f) := a_d(T)^{-1} \{ \langle X_T^{\varrho}, f \rangle - E^{\varrho} \langle X_T^{\varrho}, f \rangle \}, \quad (1.4)$$

where

$$a_d(T) = T^{1/2}, \quad d \geq 4. \quad (1.5)$$

The main result of this paper is the following.

Theorem 1.2 (Quenched CLT) *Assume $d \geq 4$ and $f \in C_p^+(\mathbb{R}^d)$. Then, for P a.e. ϱ , $Z_T^{\varrho}(f) \Rightarrow \xi(f)$ in distribution under the law P^{ϱ} as $T \rightarrow \infty$, where $\xi(f)$ is a centered Gaussian variable with variance*

$$\text{var}(\xi(f)) = \langle \lambda, fGf \rangle / 2.$$

Remarks

1. As noted above, for standard SBM in the critical dimension $d = 4$, it follows from [I86, Remark 6.1] that the occupation measure CLT norming is $(t \log t)^{1/2}$. Thus, even quenched, the random immigration has a smoothing effect on the fluctuations.
2. In [H05], the fluctuation $b_d(T)^{-1}(P^{\varrho} \langle X_T^{\varrho}, f \rangle - P^{\varrho} \langle X_T^{\varrho}, f \rangle)$ between the quenched and the annealed mean is considered. It is shown there that the choice

$$b_d(T) = \begin{cases} T^{(6-d)/4}, & 3 \leq d \leq 5, \\ (\log T)^{1/2}, & d = 6, \\ 1, & d \geq 7, \end{cases}$$

leads to non-degenerate fluctuations. Comparing Theorems 1.1 and 1.2, one sees that in dimension $d = 4$, the annealed fluctuations consist of quenched fluctuations (around the quenched mean) and of fluctuations of the quenched mean, and both contribute to the annealed variance. This is not the case for $d \geq 5$: the fluctuations of the quenched mean are of lower order and wash out in the annealed CLT.

3. For $d = 3$, an easy adaptation of our methods shows that the statement of Theorem 1.2 remains true with an almost sure statement being replaced by a statement in probability, that is $P^{\varrho}(Z_T^{\varrho}(f) > x)$ converges in probability, as $T \rightarrow \infty$, to $P(\xi(f) > x)$ for all x . Combined with the results in [H05], one concludes that for $d = 3$, the quenched fluctuations around the quenched mean are of lower order than the fluctuations of the quenched mean. Together with Theorems 1.1 and 1.2, this gives a fairly complete description of the CLT in all dimensions $d \geq 3$.
4. A functional version of Theorem 1.2 can be derived by using similar ideas. We prefer to bring here the shorter proof for the standard CLT.

2 Proof of Theorem 1.2

Set $d \geq 4$ and $f_t := a_d(t)^{-1}f$ with $f \in C_p^+(\mathbb{R}^d)$. The nonnegative mild solution of equation (1.2) is

$$v_t(r, x) = S_r f_t(x) - \int_0^r S_{r-h} v_t(h, \cdot)^2(x) dh, \quad 0 \leq r \leq t. \quad (2.1)$$

From equations (1.3) and (2.1), it follows that

$$E^\varrho \langle X_t^\varrho, f_t \rangle = \langle \lambda, S_t f_t \rangle - \int_0^t \langle \varrho_s, S_{t-s} f_t \rangle ds.$$

Combined with (1.4), we get

$$E^\varrho \exp\{-Z_t^\varrho(f)\} = \exp\left\{\langle \lambda, \int_0^t S_s v_t^2(t-s, \cdot) ds \rangle + \int_0^t \langle \varrho_s, \int_0^{t-s} S_r v_t^2(t-s-r, \cdot) dr \rangle ds\right\}. \quad (2.2)$$

The proof of Theorem 1.2 builds upon the following two propositions. The proof of Proposition 2.2 will take up most of our effort in the sequel.

Proposition 2.1 *With the above notation,*

$$\langle \lambda, \int_0^t S_s v_t^2(t-s, \cdot) ds \rangle \xrightarrow{t \rightarrow \infty} 0.$$

Set $g_t(u, x) = \int_0^u S_r v_t^2(u-r, \cdot)(x) dr = \int_0^u S_{u-r} v_t^2(r, \cdot)(x) dr$.

Proposition 2.2 *For P -a.e. ϱ ,*

$$\Gamma(t) := \int_0^t \langle \varrho_s, g_t(t-s) \rangle ds \xrightarrow{t \rightarrow \infty} \langle \lambda, fGf \rangle / 2.$$

Proof of Theorem 1.2 The theorem is an immediate consequence of (2.2), Proposition 2.1 and Proposition 2.2. \square

Proof of Proposition 2.1 A direct computation shows that, for any $d \geq 3$,

$$\int_0^\infty ds \langle \lambda, fS_{2s}f \rangle < \infty. \quad (2.3)$$

From (2.1), it follows that

$$\langle \lambda, \int_0^t S_s v_t^2(t-s, \cdot) ds \rangle = \langle \lambda, \int_0^t v_t^2(t-s, \cdot) ds \rangle \leq \int_0^t \langle \lambda, (S_{t-s} f_t)^2 \rangle ds = t^{-1} \int_0^t ds \langle \lambda, fS_{2s}f \rangle.$$

Using (2.3), the result follows. \square

Proof of Proposition 2.2 We recall from [I86, Theorem 3.2] that for any $C_p(\mathbb{R}^d)^+$ -valued continuous path $F(s)$, the Laplace transform $\int_0^t \langle \varrho_s, F(t-s) \rangle ds$ is given by

$$E \exp\left\{-\theta \int_0^t \langle \varrho_s, F(t-s) \rangle ds\right\} = \exp\{-\langle \lambda, u(t, \theta; \cdot) \rangle\}, \quad \theta > 0, \quad (2.4)$$

where $u(s, \theta; x)$ is the nonnegative solution of the following mild equation

$$u(s, \theta; x) = \theta \int_0^s S_{s-r} F(r)(x) dr - \int_0^s S_{s-r} u^2(r, \theta)(x) dr, \quad 0 \leq s \leq t. \quad (2.5)$$

(In fact, (2.4) and (2.5) hold true for $|\theta| < c$ for c a small enough constant, see [H03].) Differentiating with respect to θ in (2.4) and (2.5), we obtain

$$E \left[\int_0^t \langle \varrho_s, F(t-s) \rangle ds \right] = \int_0^t ds \int_0^s \langle \lambda, v^2(r, \cdot) \rangle dr, \quad (2.6)$$

and

$$\text{var} \left[\int_0^t \langle \varrho_s, F(t-s) \rangle ds \right] = 2 \int_0^t \langle \lambda, \left(\int_0^s S_{s-r} F(r) dr \right)^2 \rangle ds. \quad (2.7)$$

In the sequel, we let A denote a constant whose value may change from line to line and which may depend on the dimension and on f , but not on s, t, x , etc. Let us recall the useful estimate

$$\|S_s f\| \leq A \cdot (1 \wedge s^{-d/2}), \quad (2.8)$$

where $\|\cdot\|$ denotes the supremum norm.

Lemma 2.1 *Let $F(s) = g_t(s)$. Then,*

$$E \left[\int_0^t \langle \varrho_s, g_t(t-s) \rangle ds \right] \xrightarrow{t \rightarrow \infty} \langle \lambda, fGf \rangle / 2.$$

Proof. Note that

$$\begin{aligned} E \left[\int_0^t \langle \varrho_s, g_t(t-s) \rangle ds \right] &= \int_0^t ds \int_0^s \langle \lambda, v_t^2(r, \cdot) \rangle dr \\ &= \int_0^t ds \int_0^s \langle \lambda, (S_r f_t)^2 \rangle dr - \int_0^t ds \int_0^s \langle \lambda, (S_r f_t)^2 - v_t^2(r, \cdot) \rangle dr. \end{aligned}$$

One has

$$\int_0^t ds \int_0^s \langle \lambda, (S_r f_t)^2 \rangle dr = t^{-1} \int_0^t ds \int_0^s \langle \lambda, f S_{2s} f \rangle dr \xrightarrow{t \rightarrow \infty} \langle \lambda, fGf \rangle / 2.$$

On the other hand, by (2.1),

$$\begin{aligned} \int_0^t ds \int_0^s \langle \lambda, (S_r f_t)^2 - v_t^2(r, \cdot) \rangle dr &\leq 2 \int_0^t ds \int_0^s \langle \lambda, S_r f_t \cdot \int_0^r S_{r-h} v_t^2(h) dh \rangle dr \\ &\leq 2 \int_0^t ds \int_0^s \langle \lambda, S_r f_t \cdot \int_0^r S_{r-h} (S_h f_t)^2 dh \rangle dr \\ &\leq A t^{-3/2} \int_0^t ds \int_0^s \langle \lambda, (S_r f)^2 \rangle dr \cdot \int_0^t (1 \wedge h^{-d/2}) dh \\ &= A t^{-3/2} \int_0^t ds \int_0^s \langle \lambda, f S_{2r} f \rangle dr \cdot \int_0^t (1 \wedge h^{-d/2}) dh \\ &\leq \frac{A}{\sqrt{t}} \cdot \frac{1}{t} \int_0^t ds \int_0^\infty \langle \lambda, f S_{2r} f \rangle dr \end{aligned}$$

which goes to 0 when $d \geq 3$ as $t \rightarrow \infty$ due to (2.3); here, we used (2.8) at the third inequality. Substituting in (2.6), the lemma follows. \square

We return to the proof of Proposition 2.2. In view of Lemma 2.1, it is enough to prove that $\Gamma(t) - E\Gamma(t) \rightarrow 0$, as $t \rightarrow \infty$, i.e.,

$$\int_0^t \langle \varrho_s, g_t(t-s) \rangle ds - E \left[\int_0^t \langle \varrho_s, g_t(t-s) \rangle ds \right] \longrightarrow 0, \quad P \text{ a.s.} \varrho. \quad (2.9)$$

For any $n \leq t_1 \leq t \leq n+1$, let $\delta = t - t_1$, and set $\Delta\Gamma(t_1, t) := \Gamma(t) - \Gamma(t_1)$. Write $\Delta\Gamma(t_1, t)$ as the sum of four terms

$$\Delta\Gamma(t_1, t) = \Delta\Gamma_1(t_1, t) + \Delta\Gamma_2(t_1, t) + \Delta\Gamma_3(t_1, t) + \Delta\Gamma_4(t_1, t), \quad (2.10)$$

where

$$\begin{aligned} \Delta\Gamma_1(t_1, t) &:= \int_{t_1}^t \left\langle \varrho_r, \int_0^{t-r} S_{t-r-h} v_t^2(h) dh \right\rangle dr, \\ \Delta\Gamma_2(t_1, t) &:= \int_0^{t_1} \left\langle \varrho_r, \int_{t_1-r}^{t-r} S_{t-r-h} v_t^2(h) dh \right\rangle dr, \\ \Delta\Gamma_3(t_1, t) &:= \int_0^{t_1} \left\langle \varrho_r, \int_0^{t_1-r} S_{t-r-h} [v_t^2(h) - v_{t_1}^2(h)] dh \right\rangle dr, \\ \Delta\Gamma_4(t_1, t) &= \int_0^{t_1} \left\langle \varrho_r, \int_0^{t_1-r} [S_{t-r-h} - S_{t_1-r-h}] v_{t_1}^2(h) dh \right\rangle dr \end{aligned}$$

We estimate separately the moments of centered versions of $\Delta\Gamma_i(t_1, t)$.

Lemma 2.2

$$\text{var}[\Delta\Gamma_1(t_1, t)] \leq A\delta^2 n^{-2}.$$

Proof. Recall that $\Delta\Gamma_1(t_1, t) = \int_{t_1}^t \langle \varrho_r, g_t(t-r) \rangle dr$. We have, again from [I86, Theorem 3.2], that for $\theta \geq 0$,

$$\begin{aligned} E \exp \left\{ -\theta \int_{t_1}^t \langle \varrho_r, g_t(t-r) \rangle dr \right\} &= E \exp \{ -\langle \varrho_{t_1}, u(t_1, t, \theta; \cdot) \rangle \} \\ &= \exp \{ -\langle \lambda, w(0, t_1, \theta; \cdot) \rangle \}, \end{aligned}$$

where $u(s, t, \theta; \cdot)$ is the nonnegative solution of the following mild equation

$$u(s, t, \theta; x) = \theta \int_s^t S_{s,r} g_t(t-r)(x) dr - \int_s^t S_{s,r} u^2(r, t, \theta)(x) dr, \quad t_1 \leq s \leq t,$$

and $w(s, t_1, \theta; x)$ is the nonnegative solution of the following mild equation

$$w(s, t_1, \theta; x) = S_{s,t_1} u(t_1, t, \theta; \cdot)(x) - \int_s^{t_1} S_{s,r} w^2(r, t_1, \theta)(x) dr, \quad 0 \leq s \leq t_1.$$

Obviously,

$$\text{var}[\Delta\Gamma_1(t_1, t)] = - \frac{\partial^2 \langle \lambda, w(0, t_1, \theta; \cdot) \rangle}{\partial \theta^2} \Big|_{\theta=0}.$$

Performing the differentiation and using that $u|_{\theta=0} = w|_{\theta=0} = 0$, we obtain

$$\begin{aligned}
& \text{var}[\Delta\Gamma_1(t_1, t)] \\
&= 2\langle \lambda, \int_{t_1}^t S_{0,s} [\int_s^t S_{s,r} g_t(t-r) dr]^2 ds \rangle + 2\langle \lambda, \int_0^{t_1} S_{0,s} [\int_{t_1}^t S_{s,r} g_t(t-r) dr]^2 ds \rangle \\
&\leq 2t^{-2} \left[\langle \lambda, \int_{t_1}^t S_s [\int_s^t \int_0^{t-r} S_{t-s-l} (S_l f)^2 dl dr]^2 ds \rangle + \langle \lambda, \int_0^{t_1} S_s [\int_{t_1}^t \int_0^{t-r} S_{t-s-l} (S_l f)^2 dl dr]^2 ds \rangle \right] \\
&\leq At^{-2} \left[\langle \lambda, \int_{t_1}^t [(t-s) S_{t-s} f]^2 ds \rangle + \langle \lambda, \int_0^{t_1} [\int_{t_1}^t S_{t-s} f dr]^2 ds \rangle \right] \\
&\leq An^{-2} \delta^2,
\end{aligned}$$

in which (2.8) has been used several times. \square

Lemma 2.3 *With the above notation,*

$$E[\Delta\Gamma_2(t_1, t) - E\Delta\Gamma_2(t_1, t)]^4 \leq A\delta^2 n^{-2}.$$

Proof. We have $\Delta\Gamma_2(t_1, t) = \int_0^{t_1} \langle \varrho_r, F_t(t_1 - r) \rangle dr$ where $F_t(r) := \int_r^{\delta+r} S_{\delta+r-l} v_t^2(l) dl$, then $\Delta\Gamma_2(t_1, t) = \int_0^{t_1} \langle \varrho_r, F(t_1 - r) \rangle dr$. Let

$$u^{(i)}(r, x) := \frac{\partial^i u(r, x, \theta)}{\partial \theta^i} \Big|_{\theta=0}, i = 1, 2, 3.$$

Differentiating with respect to θ in (2.4) and (2.5), and using again that $u|_{\theta=0} = 0$, we obtain

$$\begin{aligned}
& E[\Delta\Gamma_2(t_1, t) - E\Delta\Gamma_2(t_1, t)]^4 \\
&= 3 \left(\int_0^{t_1} \langle \lambda, u^{(1)}(r)^2 \rangle dr \right)^2 + 3 \int_0^{t_1} \langle \lambda, u^{(2)}(r)^2 \rangle dr + 4 \int_0^{t_1} \langle \lambda, u^{(1)}(r) u^{(3)}(r) \rangle dr \\
&:= 3I^2 + 3J + 4K,
\end{aligned} \tag{2.11}$$

where for $0 \leq r \leq t_1$,

$$\begin{aligned}
u^{(1)}(r, x) &= \int_0^r S_{r-s} g_t(s) ds = \int_0^r S_{r-s} \int_s^{\delta+s} S_{\delta+s-l} v_t^2(l) dl ds \\
&\leq At_1^{-1} \int_0^r S_{r-s} \int_s^{\delta+s} S_{\delta+s-l} (S_l f)^2 dl ds \\
&\leq At_1^{-1} \delta \cdot r S_{\delta+r} f \\
|u^{(2)}(r, x)| &= \left| -2 \int_0^r S_{r-s} u'(s)^2 ds \right| \leq At_1^{-2} \delta^2 \cdot \int_0^r S_{r-s} (s S_{\delta+s} f)^2 \\
&\leq At_1^{-2} \delta^2 \cdot r S_{\delta+r} f \\
|u^{(3)}(r, x)| &= \left| -6 \int_0^r S_{r-s} u'(s) u''(s) ds \right| \leq At_1^{-3} \delta^3 \cdot \int_0^r S_{r-s} (s S_{\delta+s} f)^2 \\
&\leq At_1^{-3} \delta^3 \cdot r S_{\delta+r} f.
\end{aligned}$$

Thus, we obtain, using that $d \geq 4$,

$$I \leq At_1^{-2}\delta^2 \cdot \int_0^{t_1} \langle \lambda, (rS_{\delta+r}f)^2 \rangle dr \leq At_1^{-2}\delta^2 \cdot \int_0^{t_1} r^2(1 \wedge (\delta+r)^{-d/2})dr \leq At_1^{-1}\delta^2.$$

Similarly, with

$$J = \int_0^{t_1} \langle \lambda, u^{(2)}(r)^2 \rangle dr \leq At_1^{-4}\delta^4 \cdot \int_0^{t_1} \langle \lambda, (rS_{\delta+r}f)^2 \rangle dr \leq At_1^{-3}\delta^4,$$

and

$$K = \int_0^{t_1} \langle \lambda, u^{(1)}(r)u^{(3)}(r) \rangle dr \leq At_1^{-4}\delta^4 \cdot \int_0^{t_1} \langle \lambda, (rS_{\delta+r}f)^2 \rangle dr \leq At_1^{-3}\delta^4.$$

Substituting in (2.11) completes the proof. \square

Lemma 2.4 *With the above notation,*

$$\text{var}[\Delta\Gamma_3(t_1, t)] \leq A\delta^2 n^{-3}.$$

Proof. We begin by considering the difference $v_t(r, x) - v_{t_1}(r, x)$. From (2.1), we have

$$v_t(r, x) - v_{t_1}(r, x) = S_r f_t(x) - S_r f_{t_1}(x) + \int_0^r S_{r-h}[v_t^2(h, \cdot) - v_{t_1}^2(h, \cdot)](x)dh. \quad (2.12)$$

A direct computation reveals that $\|S_r f_t - S_r f_{t_1}\| \leq A\delta t_1^{-3/2}(1 \wedge r^{-d/2})$. Since $v_t(r, x) \leq S_r f_t$, it follows that $\|S_{r-h}[v_t^2(h, \cdot) - v_{t_1}^2(h, \cdot)]\| \leq \|2[v_t(h, \cdot) - v_{t_1}(h, \cdot)]S_r f_{t_1}\| \leq At_1^{-1/2}(1 \wedge r^{-d/2})\|v_t(h, \cdot) - v_{t_1}(h, \cdot)\|$. Thus from (2.12) we get,

$$\|v_t(r, \cdot) - v_{t_1}(r, \cdot)\| \leq A\delta t_1^{-3/2}(1 \wedge r^{-d/2}) + At_1^{-1/2}(1 \wedge r^{-d/2}) \int_0^r \|v_t(h, \cdot) - v_{t_1}(h, \cdot)\| dh.$$

Writing $a_r = \|v_t(r, \cdot) - v_{t_1}(r, \cdot)\| \geq 0$, $b_r = A\delta t_1^{-3/2}(1 \wedge r^{-d/2}) \geq 0$ and $c_r = At_1^{-1/2}(1 \wedge r^{-d/2}) \geq 0$, we thus have

$$a_r \leq b_r + c_r \int_0^r a_s ds.$$

By a version of Gronwall's inequality,

$$a_r \leq b_r + c_r \int_0^r e^{\int_s^t c_u du} b_s ds.$$

(This can be seen by setting $z_r = \int_0^r a_s ds$ and noting that z_r satisfies the differential inequality $dz_r/dr \leq b_r + c_r z_r$, with $z_0 = 0$.) Thus,

$$\begin{aligned} & \|v_t(r, \cdot) - v_{t_1}(r, \cdot)\| \\ & \leq A\delta t_1^{-3/2}(1 \wedge r^{-d/2}) + A^2 t_1^{-1/2}(1 \wedge r^{-d/2}) \int_0^r \delta t_1^{-3/2}(1 \wedge s^{-d/2}) \exp\{At_1^{-1/2} \int_s^r (1 \wedge u^{-d/2}) du\} ds \\ & \leq A\delta t_1^{-1}(1 \wedge r^{-d/2}). \end{aligned}$$

Once more by (2.12) we have

$$\begin{aligned}
& |v_t(r, x) - v_{t_1}(r, x)| \\
& \leq |S_r f_t(x) - S_r f_{t_1}(x)| + \int_0^r \|v_t(h, \cdot) - v_{t_1}(h, \cdot)\| \cdot |S_{r-h}[v_t(h, \cdot) + v_{t_1}(h, \cdot)](x)| dh \\
& \leq \delta t_1^{-3/2} S_r f + A \delta t_1^{-3/2} S_r f \cdot \int_0^r (1 \wedge h^{-d/2}) dh \\
& \leq A \delta t_1^{-3/2} S_r f.
\end{aligned}$$

Now we can estimate the variance of $\Gamma_3(t_1, t)$. By (2.7) with $F(r) = \int_0^r S_{t-t_1+r-l}[v_t^2(l) - v_{t_1}^2(l)] dl$,

$$\begin{aligned}
\text{var}[\Delta\Gamma_3(t_1, t)] &= 2 \int_0^{t_1} \left\langle \lambda, \left[\int_0^r S_{r-h} \int_0^h S_{t-t_1+h-l}[v_t^2(l) - v_{t_1}^2(l)] dl dh \right]^2 \right\rangle dr \\
&\leq A \int_0^{t_1} \left\langle \lambda, \left[\int_0^r \int_0^h S_{\delta+r-l} |v_t(l) - v_{t_1}(l)| |v_t(l) + v_{t_1}(l)| dl dh \right]^2 \right\rangle dr \\
&\leq A \delta^2 t_1^{-4} \int_0^{t_1} r^2 [1 \wedge (r + \delta)^{-d/2}] dr \\
&\leq A \delta^2 n^{-3}.
\end{aligned}$$

This completes the proof of the lemma. \square

Before providing an estimate on the moments of $\Delta\Gamma_4(t_1, t)$, we need an a-priori simple estimate on time differences of the heat kernel $p(t, x, y) = (2\pi t)^{-d/2} \exp(-|x - y|^2/2t)$. Since we did not find a direct reference for it, we provide the proof.

Lemma 2.5 *There is a constant A such that for any $t \geq \tau > 0$, we have*

$$\sup_{0 < s \leq \tau \leq t} s^{-1} |p(t + s, x, y) - p(t, x, y)| \leq A \tau^{-1} [p(t + 2\tau, x, y) + p(t, x, y)], \quad (2.13)$$

Proof. Consider first $\tau = 1$. Let $z = |x - y|$, two cases should be considered:

Case 1: $z^2 < 2d(t + 1)$. Note that

$$|p(t + s, x, y) - p(t, x, y)| = p(t, x, y) \left| \exp \left\{ \frac{z^2 s}{2t(t + s)} \right\} \left(\frac{t}{t + s} \right)^{d/2} - 1 \right|$$

and $\exp \left\{ \frac{z^2 s}{2t(t + s)} \right\} = 1 + s R_1(s, t, z)$, $\left(\frac{t}{t + s} \right)^{d/2} = \left(1 - \frac{s}{t + s} \right)^{d/2} = 1 + s R_2(s, t, z)$, where $R_1(s, t, z)$, $R_2(s, t, z)$ are bounded by a constant when $z^2 < 2d(t + 1)$, $0 < s \leq 1 \leq t$. Thus, we get

$$\sup_{0 < s \leq 1} \sup_{z^2 < 2d(t+1)} s^{-1} |p(t + s, x, y) - p(t, x, y)| \leq A p(t, x, y). \quad (2.14)$$

Case 2: $z^2 \geq 2d(t + 1)$. Since $\frac{\partial}{\partial t} p(t, z) = p(t, z) \left[-\frac{d}{2t} + \frac{z^2}{2t^2} \right]$,

$$\begin{aligned}
|p(t + s, z) - p(t, z)| &= \left| \int_0^s p(t + u, z) \left[-\frac{d}{2(t + u)} + \frac{z^2}{2(t + u)^2} \right] du \right| \\
&\leq \int_0^s p(t + u, z) \frac{z^2}{2(t + u)^2} du,
\end{aligned}$$

where the inequality uses that $\left| -\frac{d}{2(t+u)} + \frac{z^2}{2(t+u)^2} \right| \leq \frac{z^2}{2(t+u)^2}$ when $z^2 \geq 2d(t+1)$. But

$$\begin{aligned} \frac{z^2}{2(t+u)^2} p(t+u, z) &= p(t+2, z) \left(\frac{t+2}{t+u} \right)^{d/2} \frac{z^2}{(t+u)^2} \exp \left\{ -\frac{z^2}{2} \left[\frac{2-u}{(t+u)(t+2)} \right] \right\} \\ &\leq A \cdot p(t+2, z) \frac{z^2}{(t+u)^2} \exp \left\{ -\frac{z^2}{2} \left[\frac{2-u}{(t+u)(t+2)} \right] \right\}, \end{aligned}$$

(note that $0 < u \leq s \leq 1 \leq t$) and

$$\sup_{z^2 \geq 2d(t+1)} \frac{z^2}{(t+u)^2} \exp \left\{ -\frac{z^2}{2} \left[\frac{2-u}{(t+u)(t+2)} \right] \right\} < \infty.$$

So

$$\sup_{0 < s \leq 1} \sup_{z^2 \geq 2d(t+1)} s^{-1} |p(t+s, x, y) - p(t, x, y)| \leq Ap(t+2, x, y). \quad (2.15)$$

Combining (2.14) and (2.15) we obtain (2.13) when $\tau = 1$. For general $\tau > 0$, we use the scaling properties of $p(t, z)$. We have

$$\begin{aligned} s^{-1} |p(t+s, z) - p(t, z)| &= \tau^{-d/2} s^{-1} \left| p(\tau^{-1}(t+s), \tau^{-1/2}z) - p(\tau^{-1}t, \tau^{-1/2}z) \right| \\ &= \tau^{-d/2} \tau^{-1} \left[\tau s^{-1} \left| p(\tau^{-1}(t+s), \tau^{-1/2}z) - p(\tau^{-1}t, \tau^{-1/2}z) \right| \right] \\ &\leq A \tau^{-d/2} \tau^{-1} \left[p(\tau^{-1}t+2, \tau^{-1/2}z) + p(\tau^{-1}t, \tau^{-1/2}z) \right] \\ &\leq A \tau^{-1} [p(t+2\tau), z] + p(t, z), \end{aligned}$$

where the third inequality follows from the case already considered because $0 < \tau^{-1}s \leq 1 \leq \tau^{-1}t$. This complete the proof. \square

Lemma 2.6 *With the notation above, we have*

$$E[\Delta\Gamma_4(t_1, t) - E\Delta\Gamma_4(t_1, t)]^4 \leq A\delta^2 n^{-2}$$

Proof. The formula for the fourth moment of $\Delta\Gamma_4(t_1, t)$ is as in the proof of Lemma 2.3, except that the function $F_t(r)$ is replaced by the function but now $F_{t_1}(r) := \int_0^r [S_{\delta+r-l} - S_{r-l}] v_{t_1}^2(l) dl$, and $\Delta\Gamma_4(t_1, t) = \int_0^{t_1} \langle \lambda, F_{t_1}(r) \rangle dr$. Recall that $\delta = t - t_1$, we obtain for $0 \leq r \leq t_1$,

$$\begin{aligned} |u^{(1)}(r, x)| &= \left| \int_0^r S_{r-s} F_{t_1}(s) ds \right| = \left| \int_0^r S_{r-s} \int_0^s [S_{\delta+s-l} - S_{s-l}] v_{t_1}^2(l) dl ds \right| \\ &\leq \int_0^r l | [S_{\delta+l} - S_l] v_{t_1}^2(r-l) | dl \\ &\leq \int_0^\delta l [S_{\delta+l} + S_l] (S_{(r-l)} f_{t_1})^2 dl + \int_\delta^r \delta [S_{3l} + S_l] (S_{(r-l)} f_{t_1})^2 dl \\ &\leq A\delta t_1^{-1} \left[\delta (S_{\delta+r} + S_r) f + \int_\delta^r (S_{r+2l} + S_r) f \cdot (1 \wedge (r-l)^{-d/2}) dl \right], \end{aligned}$$

where the third step is from Lemma 2.5 (with $\tau = t = l, s = \delta$ there). By similar calculation we get

$$\begin{aligned}
|u^{(2)}(r, x)| &= \left| -2 \int_0^r S_{r-s} u^{(1)}(s)^2 ds \right| \\
&\leq At_1^{-2} \delta^2 \cdot \left[\delta(S_{\delta+r} + S_r) f + \int_0^r ds \int_\delta^s (S_{r+2l} + S_r) f \cdot (1 \wedge (s-l)^{-d/2}) dl \right], \\
|u^{(3)}(r, x)| &= \left| -6 \int_0^r S_{r-s} u^{(1)}(s) u^{(2)}(s) ds \right| \\
&\leq At_1^{-3} \delta^3 \cdot \left[\delta(2r^{3/2} + \delta)(S_{\delta+r} + S_r) f + r^{1/2} \int_0^r ds \int_\delta^s (S_{r+2l} + S_r) f \cdot (1 \wedge (s-l)^{-d/2}) dl \right],
\end{aligned}$$

and the estimate (2.8) was used many times. Then

$$\begin{aligned}
I &= \int_0^{t_1} \langle \lambda, u^{(1)}(r)^2 \rangle dr \\
&\leq A\delta^2 t_1^{-2} \int_0^{t_1} \left\langle \lambda, \left[\delta(S_{\delta+r} + S_r) f + \int_\delta^r (S_{r+2l} + S_r) f \cdot (1 \wedge (r-l)^{-d/2}) dl \right]^2 \right\rangle \\
&\leq A\delta^2 t_1^{-1},
\end{aligned}$$

and $J = 3 \int_0^{t_1} \langle \lambda, u^{(2)}(r)^2 \rangle dr \leq A\delta^4 t_1^{-2}$, $K = \int_0^{t_1} \langle \lambda, u^{(1)}(r) u^{(3)}(r) \rangle dr \leq A\delta^4 t_1^{-2}$. So

$$E[\Delta\Gamma_4(t_1, t) - E\Delta\Gamma_4(t_1, t)]^4 = 3I^2 + 3J + 4K \leq A\delta^4 n^{-2},$$

which completes the proof. \square

We return to the proof of Proposition 2.2. Let $\bar{\Gamma}(t) := \Gamma(t) - P\Gamma(t)$ denote the centered $\Gamma(t)$, and define $\Delta\bar{\Gamma}_i$ similarly. For any $\varepsilon > 0$,

$$\begin{aligned}
&P\left(\max_{n^2 \leq t \leq (n+1)^2} |\bar{\Gamma}(t) - \bar{\Gamma}(n^2)| > \varepsilon\right) \\
&\leq \sum_{k=1}^{\infty} P\left(\max_{0 \leq j \leq n2^k} \left| \bar{\Gamma}(n^2 + 2^{-k}(j+1)) - \bar{\Gamma}(n^2 + 2^{-k}j) \right| > \varepsilon k^{-2}/2\right) \\
&= \sum_{k=1}^{\infty} P\left(\max_{0 \leq j \leq n2^k} \left| \Delta\bar{\Gamma}(n^2 + 2^{-k}j, n^2 + 2^{-k}(j+1)) \right| > \frac{\varepsilon}{2k^2}\right) \\
&\leq \sum_{k=1}^{\infty} \sum_{i=1}^4 n2^k \max_{0 \leq j \leq n2^k} P\left(\left| \Delta\bar{\Gamma}_i(n^2 + 2^{-k}j, n^2 + 2^{-k}(j+1)) \right| > \frac{\varepsilon}{8k^2}\right).
\end{aligned}$$

By Chebyshev's inequality and Lemmas 2.2 and 2.4, for $i = 1, 3$,

$$P\left(\left| \Delta\bar{\Gamma}_i(n^2 + 2^{-k}j, n^2 + 2^{-k}(j+1)) \right| > \frac{\varepsilon}{8k^2}\right) \leq 64A\varepsilon^{-2} k^4 2^{-2k} n^{-4}.$$

Similarly, using Lemmas 2.3 and 2.6, we obtain for $i = 2, 4$,

$$P\left(\left| \Delta\bar{\Gamma}_i(n^2 + 2^{-k}j, n^2 + 2^{-k}(j+1)) \right| > \frac{\varepsilon}{8k^2}\right) \leq 4096A\varepsilon^{-4} k^8 2^{-2k} n^{-4}.$$

Thus, adjusting the value of A ,

$$P\left(\max_{n^2 \leq t \leq (n+1)^2} |\bar{\Gamma}(t) - \bar{\Gamma}(n^2)| > \varepsilon\right) \leq A\varepsilon^{-4}n^{-3} \sum_{k=1}^{\infty} k^8 2^{-k} \leq A\varepsilon^{-4}n^{-3}.$$

By the Borel-Cantelli Lemma, we get $\max_{n^2 \leq t \leq (n+1)^2} |\bar{\Gamma}(t) - \bar{\Gamma}(n^2)| \rightarrow 0$, $P - a.s.$. Thus, the proposition follows once we prove that

$$\bar{\Gamma}(n^2) \longrightarrow 0 \quad P - a.s. \quad (2.16)$$

Recall that $E[\bar{\Gamma}(n^2)] = 0$, and by (2.7),

$$\begin{aligned} \text{var}[\bar{\Gamma}(n^2)] &= \text{var} \left[\int_0^{n^2} \langle \varrho_s, g_{n^2}(n^2 - s) \rangle ds \right] = 2 \int_0^{n^2} \langle \lambda, \left(\int_0^s S_{s-r} g_{n^2}(r) dr \right)^2 \rangle ds \\ &\leq 2 \int_0^{n^2} \langle \lambda, \left[\int_0^s S_{s-r} \left(\int_0^r S_{r-l} (S_l f_{n^2})^2 dl \right) dr \right]^2 \rangle ds \\ &\leq A \cdot n^{-4} \int_0^{n^2} \langle \lambda, [s S_s f \cdot \int_0^\infty (1 \wedge l^{-d/2}) dl]^2 \rangle ds \\ &\leq A \cdot n^{-4} \int_0^{n^2} s^2 (1 \wedge s^{-d/2}) ds \\ &\leq A \cdot n^{-2}. \end{aligned}$$

Thus for any $\varepsilon > 0$,

$$\sum_{n=1}^{\infty} P[|\bar{\Gamma}(n^2)| > \varepsilon] \leq \sum_{n=1}^{\infty} \varepsilon^{-2} n^{-2} < \infty,$$

and (2.16) follows by the Borel-Cantelli Lemma. \square

Acknowledgment This work was done during a visit of Wenming Hong to the University of Minnesota. He would like to thank the Department of Mathematics, University of Minnesota for its hospitality during this visit.

References

- [D93] Dawson, D.A., Measure-valued Markov processes, In: *Lect. Notes. Math.* 1541, 1-260 (1993), Springer-Verlag.
- [DGL02] Dawson, D.A., Gorostiza, L.G., Li, Z.H., Non-local branching superprocesses and some related models, *Acta Applicandae Mathematicae*, 74 (2002), 93–112.
- [H02] Hong, W.M., Longtime behavior for the occupation time of super-Brownian motion with random immigration, *Stochastic Process. Appl.* 102 (2002), 43–62.
- [H03] Hong, W.M., Large deviations for the super-Brownian motion with super-Brownian immigration, *Journal of Theoretical Probability*, 16 (2003), 899-922.
- [H05] Hong, W.M., Quenched mean limit theorems for the super-Brownian motion with super-Brownian immigration, *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 8 (2005), 383-396.

- [HL99] Hong, W.M. and Li, Z.H., A central limit theorem for the super-Brownian motion with super-Brownian immigration, *J. Appl. Probab.* 36 (1999), 1218-1224.
- [I86] Iscoe, I., A weighted occupation time for a class of measure-valued critical branching Brownian motion, *Probab. Th. Rel. Fields* 71 (1986), 85-116.
- [I86a] Iscoe, I., Ergodic theory and a local occupation time for measure -valued critical branching Brownian motion, *Stochastics* 18 (1986), 197–243.
- [IL93] Iscoe, I., Lee, T.Y., Large deviations for occupation times of measure-valued branching Brownian motions, *Stoch. Stoch. Rep.* 45 (1993), 177-209.
- [L93] Lee, T.Y., Some limit theorems for super-Brownian motion and semilinear differential equations. *Ann. Probab.* 21 (1993), 979-995.
- [LR95] Lee, T.Y. and Remillard, B, Large deviation for three dimensional super-Brownian motion. *Ann. Probab.* 23 (1995), 1755-1771.
- [P02] Perkins, E. A., Dawson-Watanabe superprocesses and measure-valued diffusions, *Lecture Notes Math.* 1781 (2002), 132–318, Springer-Verlag.
- [RS05] Rassoul-Agha, F. and Seppäläinen, T., Ballistic random walk in a random environment with a forbidden direction, preprint (2005).
- [Ze04] Zeitouni, O., Random walks in random environment, *Lecture notes in Math.* 1837 (2004), 193–312, Springer.
- [Zh05] Zhang, M., Functional central limit theorems for the super-Brownian motion with super-Brownian immigration, *Journal of Theoretical Probability*, 18 (2005), 665-685.