

Graph coloring (homework)

The chromatic number of a graph is the minimum number of colors that suffice for a vertex coloring (in which endpoints of an edge receive different colors). This problem is NP-hard. However, this does not preclude the existence of interesting exponential time algorithms, and some of these algorithms are discussed in the lecture.

Homework assignment.

Given a graph $G(V, E)$, a *maximal independent set* is an independent set $S \subset V$ that cannot be extended by additional vertices (every vertex $v \notin S$ has some neighbor in S). Throughout we use the convention that n denotes $|V|$, and that O^* notation suppresses multiplicative terms that are polynomial in n . For example, $n^3 2^n$ is expressed as $O^*(2^n)$.

1. Give a polynomial time algorithm that given as input a graph G , outputs a maximal independent set.
2. Let $i(G)$ denote the number of maximal independent sets in the graph G . Give an algorithm that outputs *all* maximal independent sets in G , whose running time is proportional to the size of the output, namely, $O^*(i(G))$.
3. Prove that for every graph, $i(G) \leq 3^{n/2}$. (Better bounds are known, but not required in this question.)
4. For every p , show a graph on $n = 3p$ vertices (need not be connected) for which $i(G) = 3^p$, and a connected graph on $n = 3p + 1$ vertices for which $i(G) > 3^p$.
5. It is known that $i(G) \leq 3^{n/3}$ [J. W. Moon, L. Moser: On cliques in graphs. *Israel Journal of Mathematics* 3: 23-28 (1965)], and in this question you may use this fact without proof. Give a 3-coloring algorithm that runs in time $O^*(3^{n/3}) \leq O^*(1.45^n)$.