

UNIQUENESS OF SHALIKA FUNCTIONALS (THE ARCHIMEDEAN CASE)

AVRAHAM AIZENBUD, DMITRY GOUREVITCH, AND HERVÉ JACQUET

ABSTRACT. Let F be either \mathbb{R} or \mathbb{C} . Let (π, V) be an irreducible admissible smooth Fréchet representation of $GL_{2n}(F)$. A Shalika functional $\phi : V \rightarrow \mathbb{C}$ is a continuous linear functional such that for any $g \in GL_n(F)$, $A \in \text{Mat}_{n \times n}(F)$ and $v \in V$ we have

$$\phi \left[\pi \begin{pmatrix} g & A \\ 0 & g \end{pmatrix} v \right] = \exp(2\pi i \text{Re}(\text{Tr}(g^{-1}A))) \phi(v).$$

In this paper we prove that the space of Shalika functionals on V is at most one dimensional. For non-Archimedean F (of characteristic zero) this theorem was proven in [JR96].

CONTENTS

1.	Introduction	1
1.1.	Structure of the proof	2
1.2.	Structure of the paper	2
1.3.	Acknowledgements	2
2.	Preliminaries and notation	2
2.1.	Notation	2
2.2.	Admissible representations	3
3.	Integration of Shalika functionals	3
4.	Properties of $L_{\lambda, v}$	5
5.	Uniqueness of Shalika functionals	8
	References	9

1. INTRODUCTION

Let F be either \mathbb{R} or \mathbb{C} . Let (π, V) be an admissible smooth Fréchet representation of $GL_{2n}(F)$. We assume that V is the canonical completion of an irreducible Harish-Chandra (\mathfrak{g}, K) -module in the sense of Casselman-Wallach (see e.g. [Wal92], chapter 11). A **Shalika functional** $\phi : V \rightarrow \mathbb{C}$ is a continuous linear functional such that for any $g \in GL_n(F)$, $A \in \text{Mat}_{n \times n}(F)$ and $v \in V$ we have

$$\phi \left[\pi \begin{pmatrix} g & A \\ 0 & g \end{pmatrix} v \right] = \exp(2\pi i \text{Re}(\text{Tr}(g^{-1}A))) \phi(v).$$

In this paper we prove the following theorem.

Theorem 1.1. *Let (π, V) be an irreducible admissible smooth Fréchet representation of $GL_{2n}(F)$. Then the space of Shalika functionals on V is at most one dimensional.*

For non-Archimedean F (of characteristic zero) this theorem was proven in [JR96]. The proof in [JR96] is based on the fact that $(GL_{2n}(F), GL_n(F) \times GL_n(F))$ is a Gelfand pair, which was also proven in [JR96], and the method of [FJ93, Section 3] of integration of Shalika functionals.

Date: May 21, 2009.

Key words and phrases. Multiplicity one, Gelfand pair, Shalika functional, uniqueness of linear periods.

MSC Classes: 22E45.

In the Archimedean case those two ingredients also exist. Namely, [FJ93, Section 3] is valid also in the Archimedean case, and the fact that $(\mathrm{GL}_{2n}(F), \mathrm{GL}_n(F) \times \mathrm{GL}_n(F))$ is a Gelfand pair is proven in [AG08b].

The proof that we present here is similar to the proof in [JR96]. The main difference is that we have to prove the continuity of a certain linear form.

1.1. Structure of the proof.

We construct a linear map from the space of Shalika functionals to the space of linear periods (linear functionals on V that are invariant by $\mathrm{GL}_n(F) \times \mathrm{GL}_n(F)$) and prove that the map is injective. Hence the uniqueness of the linear periods implies uniqueness of the Shalika functionals. The uniqueness of linear periods, i.e. the fact that $(\mathrm{GL}_{2n}(F), \mathrm{GL}_n(F) \times \mathrm{GL}_n(F))$ is a Gelfand pair, is proven in [AG08b].

1.2. Structure of the paper.

In §2 we fix notation and terminology. In §3 we describe a way of obtaining a linear period from a Shalika functional by integration, as in [FJ93, Section 3]. In §4 we investigate the properties of the obtained period. In §5 we explain how this implies the uniqueness of Shalika functionals.

1.3. Acknowledgements.

Aizenbud and Gourevitch thank **Josef Bernstein**, **Wee Teck Gan** and **Binyong Sun** for useful remarks.

Aizenbud and Gourevitch were partially supported by a BSF grant, a GIF grant, and an ISF Center of excellency grant.

2. PRELIMINARIES AND NOTATION

2.1. Notation.

- Henceforth we fix an Archimedean field F (i.e. F is \mathbb{R} or \mathbb{C}).
- For a group G acting on a vector space V we denote by V^G the space of G -invariant vectors in V . For a character χ of G we denote by $V^{G,\chi}$ the space of (G, χ) -equivariant vectors in V .
- For a smooth real algebraic variety M we denote by $\mathcal{S}(M)$ the space of Schwartz functions on M , i.e. the space of smooth functions that are rapidly decreasing as well as all their derivatives. For precise definition see e.g. [AG08a].
- We fix a natural number n and denote $G := \mathrm{GL}_{2n}(F)$.
- We fix a norm on G by

$$\|g\| := \sum_{1 \leq i, j \leq 2n} |g_{ij}|^2 + \sum_{1 \leq i, j \leq 2n} |(g^{-1})_{ij}|^2.$$

- We denote

$$G_1 := \left\{ \begin{pmatrix} g & 0 \\ 0 & Id \end{pmatrix} \mid g \in \mathrm{GL}_n(F) \right\} \subset G$$

- We denote by $\nu : \mathrm{GL}_n(F) \rightarrow G_1$ the isomorphism defined by

$$\nu(g) := \begin{pmatrix} g & 0 \\ 0 & Id \end{pmatrix}.$$

Note that for any $X \in \mathrm{Mat}(n \times n, F)$, $d\nu(X) = \begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix}$.

- We denote

$$H := \left\{ \begin{pmatrix} g & 0 \\ 0 & g \end{pmatrix} \mid g \in \mathrm{GL}_n(F) \right\} \subset G$$

- We denote

$$U := \left\{ \begin{pmatrix} Id & A \\ 0 & Id \end{pmatrix} \mid A \in \mathrm{Mat}_{n \times n}(F) \right\} \subset G$$

- We denote by $\mu : \text{Mat}(n \times n, F) \rightarrow U$ the isomorphism defined by

$$\mu(A) := \begin{pmatrix} Id & A \\ 0 & Id \end{pmatrix}.$$

Note that for any $X \in \text{Mat}(n \times n, F)$, $d\mu(X) = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$.

- We denote by $\tau : U \rightarrow F$ the homomorphism given by

$$\tau(\mu(A)) := \text{Tr}(A).$$

- We let ψ be the additive character of F defined by $\psi(x) := e^{2\pi i \text{Re } x}$. We define an homomorphism $\Psi : U \rightarrow F^\times$ by

$$\Psi := \psi \circ \tau.$$

We extend Ψ to an homomorphism $\Psi : HU \rightarrow F^\times$ trivial on H .

- We denote by K the standard maximal compact subgroup of G . Thus $K = O(2n)$ if $F = \mathbb{R}$ and $K = U(2n)$ if $F = \mathbb{C}$.

2.2. Admissible representations.

In this paper we consider admissible smooth Fréchet representations of G , i.e. smooth admissible representations (π, V) of G such that V is a Fréchet space and, for any continuous semi-norm α on V , there exist another continuous semi-norm β on V and a natural number M such that for any $g \in G$,

$$\alpha(\pi(g)v) \leq \beta(v) \|g\|^M.$$

By Casselman - Wallach theorem (see e.g. [Wal92], chapter 11), V may be regarded as the canonical model of an irreducible Harish-Chandra (\mathfrak{g}, K) -module. By Casselman embedding theorem ([Cas80]), V can be realized as a closed subspace of a principal series representation. We denote by \tilde{V} the canonical model of the contragredient Harish-Chandra (\mathfrak{g}, K) -module. It is a subspace of the topological dual V^* of V .

3. INTEGRATION OF SHALIKA FUNCTIONALS

In this section we fix:

- an irreducible admissible smooth Fréchet representation (π, V) of G
- a Shalika functional λ on V , i.e. $\lambda \in (V^*)^{HU, \Psi}$.

Theorem 3.1. *There exists $M \in \mathbb{R}$ such that for any $v \in v$ and over the region of $s \in \mathbb{C}$ with $\text{Re}(s) > M$, the integral*

$$L_{\lambda, v}(s) := \int_{g \in G_1} \lambda(\pi(g)v) |\det(g)|^{s-\frac{1}{2}} dg$$

converges absolutely and is a holomorphic function of s .

Moreover, $L_{\lambda, v}(s)$ has meromorphic continuation to the complex plane and is a holomorphic multiple of the L -function L_π of the representation π . Finally, for any $\lambda \neq 0$ there exists $v \in V$ such that $L_{\lambda, v} = L_\pi$.

In [FJ93, Proposition 3.1] this theorem is proven under the following assumption:

(*) There exists a continuous semi-norm β on V such that $|\lambda(\pi(g)v)| \leq \beta(v)$ for any $g \in G$.

This may not be true in general. However, we have the following result.

Lemma 3.2. *There exist $M > 0$ and a continuous semi-norm β on V such that $|\lambda(\pi(g)v)| \leq |\det g|^{-M} \beta(v)$ for any $g \in G_1$.*

Before proving the Lemma, we check that, with the help of this Lemma, the proof of Theorem 3.1 is still valid. Indeed, the functions $g \mapsto \lambda(\pi(g)v)$ are bounded in [FJ93] and satisfy a sharper estimate ([FJ93, Lemma 3.1]). Here they satisfy the following estimates.

Lemma 3.3. *There is a continuous semi-norm γ on V such that, for any $v \in V$,*

$$|\lambda(\pi(g)v)| \leq |\det b^{-1}a|^{-M} \gamma(v)$$

for

$$g = u \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} k$$

with $a, b \in GL(n, F)$, $u \in U$, $k \in K$. Furthermore, for any $v \in V$, there is $\Phi_v \in \mathcal{S}(\text{Mat}(n \times n, F))$ such that

$$|\lambda(\pi(g)v)| \leq \Phi(b^{-1}a) |\det b^{-1}a|^{-M},$$

for g of the above form.

Proof. For the first assertion, we have

$$\lambda(\pi(g)v) = \Psi(u) \lambda(\pi(\nu(b^{-1}a)) \pi(k)v).$$

Hence

$$|\lambda(\pi(g)v)| \leq |\det b^{-1}a|^{-M} \beta(\pi(k)v).$$

There is another continuous semi-norm γ such that, for all $k \in K$,

$$\beta(\pi(k)v) \leq \gamma(v).$$

The first assertion follows.

For the second assertion, we go through the proof of [FJ93, Lemma 3.1] (which is the above estimate with $M = 0$) and arrive at once at the present estimate. \square

The proof of Theorem 3.1 is still valid. The only modification is that we need to check that, under our weaker assumption, two integrals in [FJ93] which depend on $s \in \mathbb{C}$, are still absolutely convergent for $\text{Re } s \gg 0$.

The first integral is integral [FJ93, 45]:

$$\int \lambda(\pi(g)v) \Phi(g) |\det g|^{s+n-\frac{1}{2}} d^\times g$$

where $\Phi \in \mathcal{S}(\text{Mat}(2n \times 2n, F))$. We write

$$g = \begin{pmatrix} a & x \\ 0 & b \end{pmatrix} k.$$

Then

$$d^\times g = |\det a|^{-n} d^\times a d^\times b dx dk.$$

By Lemma 3.3, the integral of the absolute value is bounded by

$$\int |\det a|^{\text{Re } s - M - \frac{1}{2}} |\det b|^{\text{Re } s + M + n - \frac{1}{2}} \left| \Phi \left[\begin{pmatrix} a & x \\ 0 & b \end{pmatrix} k \right] \right| d^\times a d^\times b dx dk.$$

This does converge absolutely for $\text{Re } s \gg 0$.

The second integral is integral [FJ93, 48]. It has the form

$$\int \lambda \left[\pi \begin{pmatrix} a & 0 \\ 0 & \text{Id} \end{pmatrix} \pi(x)v \right] |\det a|^{s-\frac{1}{2}} d^\times a d\mu(x)$$

where μ is the measure on $SL(2n, F)$ defined by

$$\int f(x) d\mu(x) = \int f \left[\begin{pmatrix} b^{-1} & 0 \\ 0 & \text{Id} \end{pmatrix} \begin{pmatrix} \text{Id} & u \\ 0 & \text{Id} \end{pmatrix} \begin{pmatrix} \text{Id} & 0 \\ 0 & b \end{pmatrix} k \right] \Upsilon(u, b^{-1}, b; k) |\det b|^n d^\times b du dk$$

In this formula k is integrated over $K' = K \cap SL(2n, F)$ and the function Υ is in $\mathcal{S}(\text{Mat}(n \times n, F)^3 \times K')$.

The integral of the absolute value of the integrand is bounded by

$$\int \Phi_v(ab^{-2}) |\det ab^{-2}|^{-M} |\Upsilon|(u, b^{-1}, b; k) |\det b|^n d^\times b du dk |\det a|^{\text{Re } s - \frac{1}{2}} d^\times a.$$

After changing a to ab^2 , the integral decomposes into a product:

$$\int \Phi_v(a) |\det a|^{\operatorname{Re} s - M - \frac{1}{2}} d^\times a \times \int |\Upsilon|(u, b^{-1}, b; k) |\det b|^{n+2\operatorname{Re} s - 1} d^\times b d u d k.$$

The first integral converges for $\operatorname{Re} s \gg 0$. The second integral converges for all s .

It remains to prove Lemma 3.2. We will prove the following more general lemma.

Lemma 3.4. *There exists $M_0 > 0$ such that, for any polynomial P on the real vector space $\operatorname{Mat}(n \times n, F)$, there exists a continuous semi-norm β_P on V such that for any $g \in \operatorname{GL}_n(F)$ we have*

$$|\lambda(\pi(\nu(g))v)| \leq \beta_P(v) \frac{1}{|P(g)|} |\det g|^{-M_0}.$$

Proof. We have

$$\lambda(\pi(\mu(X))v) = \psi(\operatorname{Tr} X) \lambda(v) \quad \forall X \in \operatorname{Mat}(n \times n, F).$$

We have then

$$\lambda(d\pi(d\mu(X))v) = 2\pi i \operatorname{Re} \operatorname{Tr}(X) \lambda(v) \quad \forall X \in \operatorname{Mat}(n \times n, F)$$

and hence

$$\lambda(\pi(\nu(g))d\pi(d\mu(X))v) = 2\pi i \operatorname{Re} \operatorname{Tr}(gX) \lambda(\pi(\nu(g))v) \quad \forall X \in \operatorname{Mat}(n \times n, F) \text{ and } g \in \operatorname{GL}_n(F).$$

Similarly, if Q is a polynomial on the real vector space $\operatorname{Mat}(n \times n, F)$, there is an element X_Q of the enveloping algebra of $\mathfrak{gl}_{2n}(F)$ such that

$$\lambda(\pi(\nu(g))d\pi(X_Q)v) = Q(g) \lambda(\pi(\nu(g))v) \quad \forall g \in \operatorname{GL}_n(F).$$

We know that there exist a continuous semi-norm β on V and a natural number M such that $|\lambda(\pi(g)v)| \leq \beta(v) \|g\|^M$ for any $g \in G$. Therefore for any $g \in \operatorname{GL}_n(F)$ we have

$$|Q(g) \lambda(\pi(\nu(g))v)| = |\lambda(\pi(\nu(g))d\pi(\mu(X_Q))v)| \leq \beta(d\pi(X_Q)v) \|\nu(g)\|^M.$$

Note that $\|\nu(g)\|^M = P_0(g) |\det g|^{-2M}$ for a suitable polynomial P_0 on the real vector space $\operatorname{Mat}(n \times n, F)$. Therefore, we have, with $M_0 = 2M$,

$$|\lambda(\pi(\nu(g))v)| \leq \beta(d\pi(X_Q)v) \frac{P_0(g)}{|Q(g)|} |\det g|^{-M_0}.$$

We may take Q of the form $Q = P_0 P$ where P is another polynomial. Since $v \mapsto \beta(d\pi(X_Q)v)$ is a continuous semi-norm the Lemma follows. \square

4. PROPERTIES OF $L_{\lambda, v}$

Theorem 4.1. *Let (π, V) be an irreducible admissible smooth Fréchet representation of G . Fix a Shalika functional $\lambda \in (V^*)^{HU, \Psi}$ and a vector $v \in V$. Then, for any polynomial p , the product $p(s)L_{\lambda, v}(s)$ is bounded at infinity on every vertical strip of finite width.*

In [FJ93, §§3.3] the following statement is proven.

Lemma 4.2. *For $\operatorname{Re}(s)$ large enough, $L_{\lambda, v}(s)$ is a finite sum of functions of the type*

$$\mathcal{L}_{u, \xi, \Phi}(s) := \int_{g \in G} \Phi(g) \xi(\pi(g)u) |\det g|^{s+n-\frac{1}{2}} dg,$$

where $\Phi \in \mathcal{S}(\operatorname{Mat}(2n \times 2n, F))$, $u \in V$, $\xi \in V^*$.

Now Theorem 4.1 follows from the following one.

Theorem 4.3. *Let (π, V) be an irreducible admissible smooth Fréchet representation of G . Let $\Phi \in \mathcal{S}(\operatorname{Mat}(2n \times 2n, F))$, $u \in V$ and $\xi \in \tilde{V}$. Then $\mathcal{L}_{u, \xi, \Phi}(s)$ has a meromorphic continuation to \mathbb{C} whose product by any polynomial is bounded at infinity on any vertical strip. The continuation is a holomorphic multiple of $L_\pi(s) = L(s, \pi)$. It satisfies the functional equation*

$$\int \widehat{\Phi}(g) \xi(\pi({}^t g^{-1})u) |\det g|^{1-s+n-\frac{1}{2}} dg = \gamma(s, \pi, \psi) \mathcal{L}_{u, \xi, \Phi}(s)$$

where

$$\gamma(s, \pi, \psi) := \varepsilon(s, \pi, \psi) \frac{L(1-s, \tilde{\pi})}{L(s, \pi)}.$$

and

$$\widehat{\Phi}(X) := \int_{\text{Mat}(2n \times 2n, F)} \Phi(Y) \psi(\text{tr}(XY^t)) dY.$$

Finally, these assertions remain true if ξ is in V^* (topological dual of V).

This theorem is proven in [GJ72] in slightly narrower generality: the vectors u and ξ are K -finite and the function Φ is the product of a Gaussian function and a polynomial. For the convenience of the reader we indicate how to extend the results of [GJ72].

We will need the following lemma.

Lemma 4.4. *Let $T \subset G$ be the torus of diagonal matrices. We will also regard T as the subset $(F^\times)^{2n}$ of F^{2n} . Let $\chi : T \rightarrow \mathbb{C}^\times$ be a multiplicative character. Let (π, V) be the corresponding representation of principal series of G . Let $v \in V$ and $\xi \in \tilde{V}$. Let Φ be a Schwartz function on $\text{Mat}(2n \times 2n, F)$.*

Then there exists a Schwartz function $\phi \in \mathcal{S}(F^{2n})$ such that

$$\int_{g \in G} \Phi(g) \xi(\pi(g)v) |\det g|^{s+n-\frac{1}{2}} dg = \int_{t \in T} \phi(t) \chi(t) |\det t|^s dt$$

for any $s \in \mathbb{C}$ such that the integral on the right converges absolutely.

Proof. Let N denote the group of upper triangular matrices with unit diagonal. Let $B = TN$ and δ_B be the module of the group B . Realize V as the space of smooth functions on G that satisfy

$$f(tg) = \chi(t) f(g) \delta_B^{1/2}(t) \text{ and } f(ug) = f(g) \text{ for any } t \in T \text{ and } u \in N.$$

Realize also \tilde{V} in the corresponding way. Then

$$\xi(\pi(g)v) = \int_{k \in K} v(kg) \xi(k) dk,$$

where K is the standard maximal compact subgroup. Now

$$\int_G \Phi(g) \xi(\pi(g)v) |\det g|^{s+\frac{n-1}{2}} dg = \int_G \int_K \Phi(g) v(kg) \xi(k) |\det g|^{s+\frac{n-1}{2}} dg dk = \int_G \int_K \Phi(k^{-1}g) v(g) \xi(k) |\det g|^{s+\frac{n-1}{2}} dg dk$$

To compute this integral we set

$$g = \begin{pmatrix} a_1 & u_{1,2} & \cdots & u_{1,2n} \\ 0 & a_2 & \cdots & u_{2,2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_{2n} \end{pmatrix} k'.$$

Then

$$dg = |a_1|^{1-2n} |a_2|^{2-2n} \cdots |a_{2n-1}|^{-1} \otimes d^\times a_i \otimes du_{i,j} dk'$$

We set

$$\phi(a_1, \dots, a_{2n}) := \int \Phi \left[k^{-1} \begin{pmatrix} a_1 & u_{1,2} & \cdots & u_{1,2n} \\ 0 & a_2 & \cdots & u_{2,2n} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & a_{2n} \end{pmatrix} k' \right] v(k') \xi(k) dk dk' \otimes du_{i,j}.$$

Clearly ϕ is a Schwartz function on F^{2n} and

$$\int_{g \in G} \Phi(g) \xi(\pi(g)v) |\det g|^{s+\frac{n-1}{2}} dg = \int_{t \in T} \phi(t) \chi(t) |\det t|^s dt$$

for any $s \in \mathbb{C}$ such that the integral on the right converges. □

Now we can prove Theorem 4.3.

Proof of Theorem 4.3. The representation (π, V) is a sub-representation of a principal series representation determined by a character χ of T and the representation $(\tilde{\pi}, \tilde{V})$ is then a quotient of the representation determined by χ^{-1} . For $u \in V$ and $\xi \in \tilde{V}$ (or ξ in the principal series determined by χ^{-1}) we have

$$\mathcal{L}_{u,\xi,\Phi}(s) = \int_{(a_1, \dots, a_{2n}) \in F^{\times 2n}} \phi(a_1, a_2, \dots, a_{2n}) \chi_1(a_1) |a_1|^s \dots \chi_{2n}(a_{2n}) |a_{2n}|^s d^\times a_1 \dots d^\times a_{2n}.$$

The right hand side extends to a meromorphic function of s and the product of this function by any polynomial is bounded at infinity in any vertical strip. Moreover, the function ϕ depends continuously on $\Phi, u \in V, \xi \in \tilde{V}$. Therefore the analytic continuation depends continuously on $\Phi, v \in V, \xi \in \tilde{V}$. By continuity, it has the properties stated in the Theorem. To extend further to the case where ξ is in the topological dual V^* we appeal to the Dixmier Malliavin Lemma ([DM78]) applied to the representation of $SL_{2n}(F)$ on $\mathcal{S}(\text{Mat}_{2n \times 2n}(F))$ defined by

$$g_1 \Phi(X) := \Phi(g_1^{-1} X)$$

Thus we may assume Φ is of the form

$$\Phi(X) = \int_{SL_{2n}(F)} \Phi_1(g_1^{-1} X) f(g_1) dg_1$$

where f_1 is a C^∞ function of compact support on $SL_{2n}(F)$. Then

$$\mathcal{L}_{u,\xi,\Phi} = \mathcal{L}_{u,\xi_1,\Phi_1}$$

where

$$\xi_1(v) := \xi(\pi(f_1)v).$$

Now ξ_1 is in \tilde{V} and our assertion follows. □

Remark 4.5. *The previous result with $\xi \in V^*$ is used without comment in [FJ93], formula (57). This is why we included a sketch of the proof.*

Theorem 4.6. *There exists $M > 0$ such that for any even integer $M' \geq 2$ and any polynomial p on \mathbb{C} , there exists a semi-norm β on V such that $|pL_{\lambda,v}|_{M'+M+i\mathbb{R}}| \leq \beta(v)$.*

First, we will prove the following lemma.

Lemma 4.7. *There exists $M > 0$ such that, for any even integer $M' \geq 2$, there exists a continuous semi-norm β on V such that $|L_{\lambda,v}(s)| \leq \beta(v)$ for $\text{Re } s = M' + M$.*

Proof. By Lemma 3.4 there exists $M_0 > 0$, and for any polynomial P on $\text{Mat}(n \times n, F)$, a continuous semi-norm β_P such that

$$\lambda(\pi(\nu(g))v) \leq \beta_P(v) \frac{1}{|P(g)|} |\det g|^{-M_0}.$$

Let $M := 1/2 + n^2 + M_0$. Let $M' \geq 2$ be an even integer and let $s \in \mathbb{C}$ with $\text{Re}(s) = M + M'$. Let

$$P(X) = |\det X|^{M'} \prod_{i,j} (1 + X_{ij} \bar{X}_{ij}).$$

Let

$$\beta(v) := \beta_P(v) \int_{X \in \text{Mat}(n \times n, F)} \frac{dX}{\prod_{i,j} (1 + X_{ij} \bar{X}_{ij})}.$$

Now

$$\begin{aligned} |L_{\lambda,v}| &= \left| \int_{GL_n(F)} \lambda(\pi(\nu(g))v) |\det g|^{s-1/2} dg \right| \leq \int_{GL_n(F)} \beta_P(v) \frac{1}{|P(g)|} |\det g|^{-M_0} |\det g|^{n^2+M'+M_0} dg = \\ & \int_{GL_n(F)} \beta_P(v) \frac{1}{|P(g)|} |\det g|^{n^2+M'} dg = \int_{X \in \text{Mat}(n \times n, F)} \beta_P(v) \frac{1}{|P(X)|} |\det X|^{M'} dX = \beta(v). \end{aligned}$$

□

Proof of Theorem 4.6. For any $g \in GL_n(F)$ we have

$$L_{\lambda, \pi(\nu(g))_v}(s) = |\det(g)|^{1/2-s} L_{\lambda, v}(s).$$

We can apply this to $g = \exp(tX)$, with $t \in \mathbb{R}$ and $X \in \text{Mat}(n \times n, F)$. We get

$$L_{\lambda, \pi(\nu(g))_v}(s) = |\det(\exp(tX))|^{1/2-s} L_{\lambda, v}(s).$$

Differentiating this identity with respect to t at $t = 0$, we get

$$L_{\lambda, d\pi(d\nu(X))_v}(s) = \left(\frac{1}{2} - s\right) c(X) L_{\lambda, v}(s),$$

where $c(X) = \text{Tr } X$ if $F = \mathbb{R}$ and $c(X) = 2 \text{Re } \text{Tr } X$ if $F = \mathbb{C}$. Similarly, for any polynomial p on \mathbb{C} there exists X_p in the universal enveloping algebra of $\mathfrak{gl}_{2n}(F)$ such that

$$L_{\lambda, d\pi(X_p)_v}(s) = p(s) L_{\lambda, v}(s).$$

The theorem follows now from Lemma 4.7. \square

Notation 4.8. Define another representation π^θ on the same space V by $\pi^\theta(g) := \pi((g^t)^{-1})$. Recall that $\pi^\theta \cong \tilde{\pi}$.

For any Shalika functional $\lambda : \pi \rightarrow \mathbb{C}$ we define $\lambda^\theta : \pi^\theta \rightarrow \mathbb{C}$ by

$$\lambda^\theta(v) := \lambda \left(\pi \left(\begin{pmatrix} 0_{nn} & Id_{nn} \\ -Id_{nn} & 0_{nn} \end{pmatrix} v \right) \right).$$

It is easy to see that λ^θ is a Shalika functional for the representation π^θ .

Theorem 4.9 ([FJ93], Proposition 3.3).

$$\gamma(s, \pi, \psi) L_{\lambda, v}^\pi(s) = L_{\lambda^\theta, v}^{\pi^\theta}(1-s).$$

Using Theorem 4.6 we obtain the following corollary.

Corollary 4.10. *There exists $N < 0$ such that for any odd integer $N' \leq -1$ and any polynomial p on \mathbb{C} there exists a semi-norm β on V such that $|pL_{\lambda, v}|_{N'+N+i\mathbb{R}}| \leq \beta(v)$.*

5. UNIQUENESS OF SHALIKA FUNCTIONALS

Theorem 5.1. *Let (π, V) an irreducible admissible representation of G . Let λ be a Shalika functional. Then the functional $L(\lambda) : V \rightarrow \mathbb{C}$ defined by*

$$L(\lambda)(v) := \frac{L_{\lambda, v}}{L_\pi} \left(\frac{1}{2} \right)$$

is continuous.

Proof. By Theorem 4.6 we choose $M > 1$ such that for any polynomial p there exists a semi-norm β on V such that $|pL_{\lambda, v}|_{M+i\mathbb{R}}| \leq \beta(v)$. By Corollary 4.10 we choose $N < 0$ such for any polynomial p there exists a semi-norm β' on V such that $|pL_{\lambda, v}|_{N+i\mathbb{R}}| \leq \beta'(v)$. Let q be a polynomial such that the multiset of poles of $1/q$ (with multiplicities) coincides with the multiset of poles of $L_\pi|_{[N, M]+i\mathbb{R}}$. Here, $[N, M] + i\mathbb{R}$ denotes the strip $N \leq \text{Re}(s) \leq M$. It is enough to show that the map $L'(\lambda)$ defined by $L'(\lambda)(v) := L_{\lambda, v} q(\frac{1}{2})$ is continuous. Now there exists a semi-norm α on V such that, for any $v \in V$,

$$|qL_{\lambda, v}|_{M+i\mathbb{R}}| \leq \alpha(v) \text{ and } |qL_{\lambda, v}|_{N+i\mathbb{R}}| \leq \alpha(v).$$

By Theorem 4.1, for any $v \in V$, there exists Δ such that $|qL_{\lambda, v}(s)| \leq \alpha(v)$ if $s \in [N, M] + i\mathbb{R}$ and $|\text{Im } s| \geq \Delta$. Now by maximal modulus principle $L'(\lambda)(v) \leq \alpha(v)$ for any $v \in V$. \square

Definition 5.2. *Let (π, V) an irreducible admissible representation of G . We define a map*

$$L : (V^*)^{HU, \Psi} \rightarrow (V^*)^{HG_1}$$

by

$$L(\lambda)(v) = \frac{L_{\lambda, v}}{L_\pi} \left(\frac{1}{2} \right)$$

Theorem 3.1 implies

Proposition 5.3. *L is a monomorphism.*

Now we use the following theorem from [AG08b].

Theorem 5.4 (see [AG08b], Theorem I). *The pair $(GL_{2n}, GL_n \times GL_n)$ is a Gelfand pair. Namely,*

$$\dim(V^*)^{GL_n(F) \times GL_n(F)} \leq 1.$$

Corollary 5.5. *Theorem 1.1 holds. Namely,*

$$\dim(V^*)^{HU, \Psi} \leq 1.$$

REFERENCES

- [AG08a] A. Aizenbud, D. Gourevitch, *Schwartz functions on Nash Manifolds*, International Mathematics Research Notices, Vol. 2008, n.5, Article ID rnm155, 37 pages. DOI: 10.1093/imrn/rnm155. See also arXiv:0704.2891 [math.AG].
- [AG08b] A. Aizenbud, D. Gourevitch, *An Archimedean analog of Jacquet - Rallis theorem*, arXiv:0803.3397v3 [math.RT].
- [Cas80] W. Casselman, *Jacquet modules for real reductive groups*, in Proceedings of the International Congress of Mathematicians (Helsinki, 1978), 557563, Acad. Sci. Fennica, Helsinki.
- [CHM00] W. Casselman; H. Hecht; D. Milićić, *Bruhat filtrations and Whittaker vectors for real groups*, The mathematical legacy of Harish-Chandra (Baltimore, MD, 1998), 151-190, Proc. Sympos. Pure Math., **68**, Amer. Math. Soc., Providence, RI, (2000).
- [DM78] Jacques Dixmier and Paul Malliavin, *Factorisations de fonctions et de vecteurs indefiniment differentiables*. Bull. Sci. Math. (2) **102** (1978), no. 4, 307330. MR MR517765
- [GJ72] R. Godement and H. Jacquet, *Zeta functions of simple algebras*, Lecture Notes in Mathematics **260**, Berlin-Heidelberg-New York (1972).
- [FJ93] Solomon Friedberg and Hervé Jacquet, *Linear periods*. J. Reine Angew. Math. **443** (1993), 91–139. MR MR1241129 (94k:11058)
- [JR96] Hervé Jacquet and Stephen Rallis, *Uniqueness of linear periods*, Compositio Math. **102** (1996), no. 1, 65–123. MR MR1394521 (97k:22025)
- [Wal88] N. Wallach, *Real Reductive groups I*, Pure and Applied Math. **132**, Academic Press, Boston, MA (1988).
- [Wal92] N. Wallach, *Real Reductive groups II*, Pure and Applied Math. **132-II**, Academic Press, Boston, MA (1992).

AVRAHAM AIZENBUD, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, THE WEIZMANN INSTITUTE OF SCIENCE POB 26, REHOVOT 76100, ISRAEL.

E-mail address: aizenr@yahoo.com

DMITRY GOUREVITCH, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, THE WEIZMANN INSTITUTE OF SCIENCE POB 26, REHOVOT 76100, ISRAEL.

E-mail address: guredim@yahoo.com

HERVÉ JACQUET, MATHEMATICS DEPARTMENT OF COLUMBIA UNIVERSITY, MC 4406, 2990 BROADWAY NEW YORK, NY 10027

E-mail address: hj@math.columbia.edu