# Linear Programming Duality

May 15, 2002

LP-duality asserts that if the optimal solution of the primal linear program below exists and is bounded, then the same holds for the dual linear program, and both optimal solutions have the same value. (Here $A_i$ denotes row $i$ and $A^j$ denotes column $j$.)

**Primal:** *minimize* $c^T x$ subject to
$A_i x \geq b_i$, $1 \leq i \leq h$
$A_i x = b_i$, $h < i \leq m$
$x_j \geq 0$, $1 \leq j \leq \ell$.

**Dual:** *maximize* $b^T y$ subject to
$(A^j)^T y \leq c_j$, $1 \leq j \leq \ell$
$(A^j)^T y = c_j$, $\ell < j \leq n$
$y_i \geq 0$, $1 \leq i \leq h$.

It may happen that neither the primal LP nor its dual are feasible, or that one is unbounded and the other is not feasible.

The duality theory has many consequences. In particular it places linear programming in NP $\bigcap$ coNP, it implies that finding a feasible solution to a linear program is essentially as difficult as finding an optimal solution, it can be used in order to prove other duality type theorems. One such theorem is Farkas' lemma that says that either $Ax = b$ has a solution with $x \geq 0$, or there is a vector $y$ such that $y^T A \geq 0$ but $y^T b < 0$. Whenever one studies a linear program, it is informative to look at its dual.

In economic theory, dual variables are interpreted as *shadow prices.*

Let $x$ and $y$ be feasible solutions to the primal and dual problem. Then they are optimal iff they satisfy the *complementary slackness* conditions. Namely $y^T(Ax - b) = 0$ and $(c^T - y^T A)x = 0$.

**Homework**

1. Prove that the dual of the dual is the primal. That is, take an LP minimization problem in general form (involving equality and inequality constraints, nonnegative and free variables), follow the syntactic rules for writing its dual, negate the objective function of the dual to get a minimization problem, write its dual, again negate the objective function, and check that you arrived at an LP equivalent to the primal.

2. Helly's theorem says that for any collection of convex bodies in $R^n$, if every $n+1$ of them intersect, then there is a point lying in the intersection of all of them. Prove Helly's theorem for the special case that the convex bodies are halfspaces. (Hint: show that if a system of inequalities $Ax \geq b$ does not have a solution, then we can select $n+1$ of the inequalities such that the resulting system does not have a solution. Construct a primal LP from this system by choosing a 0 cost vector.)