

# Randomized Algorithms 2017A – Lecture 13

## The JL Transform and Oblivious Subspace Embedding\*

Robert Krauthgamer

### 1 The JL Transform

**JL dimension reduction:** We saw the JL lemma which reduces the dimension of  $n$  points in  $\mathbb{R}^d$ . Recall that it uses a random linear map that is drawn obliviously of the data and works with high probability.

Below, we abstract its performance guarantee and ignore the implementation, because algorithms may have different tradeoffs, e.g., between the target dimension and the runtime.

Here is a good way to think about the next definition. A matrix  $S \in \mathbb{R}^{s \times n}$  which is just a linear map  $S : \mathbb{R}^n \rightarrow \mathbb{R}^s$ . It will represent dimension reduction, where a problem involving an unknown set of  $b$  points in  $\mathbb{R}^n$  is reduced to points in dimension  $s = s(n, b, \varepsilon, \delta)$ . Thus, we want  $s$  (the number of rows in  $S$ ) to be as small as possible.

Throughout, all norms are  $\ell_2$ -norms.

**Definition:** A random matrix  $S \in \mathbb{R}^{s \times n}$  is called an  $(\varepsilon, \delta, b)$ -Johnson-Lindenstrauss Transform (JLT) if

$$\forall B \subset \mathbb{R}^n, |B| \leq b, \quad \Pr_S \left[ \forall x \in B, \|Sx\| \in (1 \pm \varepsilon)\|x\| \right] \geq 1 - \delta.$$

We saw in class that a matrix of independent Gaussians (scaled appropriately) attains this guarantee, with a suitable  $s = O(\varepsilon^{-2} \log(b/\delta))$ . More precisely, we saw it only for  $b = 1$ , but general  $b$  follows easily by applying that result with smaller  $\delta' = \delta/b$  and taking a union bound over  $B$ .

Notice that the target dimension does not depend on the ambient dimension  $d$ .

We saw also another construction, with bigger target dimension  $s$ , but faster matrix-vector multiplication (back then we called it  $L = SHD$ ).

---

\*These notes summarize the material covered in class, usually skipping proofs, details, examples and so forth, and possibly adding some remarks, or pointers. The exercises are for self-practice and need not be handed in. In the interest of brevity, most references and credits were omitted.

## 2 Approximate Matrix Multiplication

**Definition:** The *Frobenius norm* of a real matrix  $A$  is defined as

$$\|A\|_F := \left(\sum_{i,j} A_{ij}^2\right)^{1/2}.$$

**Problem definition:** In *Approximate Matrix Multiplication (AMM)*, the input is  $\varepsilon > 0$  and two matrices  $A, B \in \mathbb{R}^{n \times m}$ , and the goal is to compute a matrix  $C \in \mathbb{R}^{m \times m}$  such that

$$\|A^T B - C\|_F \leq \varepsilon \|A\|_F \|B\|_F.$$

**Theorem 1:** Suppose the matrix  $S \in \mathbb{R}^{n \times s}$  is  $(\varepsilon', \delta', b')$ -JLT, where the parameters satisfy  $\varepsilon' = \Theta(\varepsilon)$ ,  $\delta' = \delta$ , and  $b' = O(m^2)$ . Then with probability at least  $1 - \delta$ , the matrix  $(SA)^T(SB)$  solves AMM.

Roughly speaking, this theorem reduces the dimension  $n$  (of the input matrices) with dimension  $s \approx O(\varepsilon^{-2} \log m)$ .

**Proof:** Was seen in class. The main idea is that for fixed  $x, y \in \mathbb{R}^n$  with  $\|x\| = \|y\| = 1$ , we have that

$$2[\langle Sx, Sy \rangle - \langle x, y \rangle] = [\|Sx\|^2 - \|x\|^2] + [\|Sy\|^2 - \|y\|^2] - [\|Sx - Sy\|^2 - \|x - y\|^2].$$

And now by the JLT guarantee, with high probability  $1 - \delta'$ , each of the three summands is bounded in absolute value.

**Remark:** The above proof bounds the error in each  $C_{i,j}$  (output entry) with high probability, but it clearly suffices to bound the expected squared error, which can be achieved with a smaller matrix  $S$  (e.g., no dependence on  $b' = O(m^2)$ ).

## 3 Oblivious Subspace Embedding

**Embedding an entire subspace:** In some situations (like regression, as we will see soon), we want a guarantee for a whole subspace, which has infinitely many points.

Observe that a linear subspace  $V \subset \mathbb{R}^n$  of dimension  $d$  can be described as the column space of  $A \in \mathbb{R}^{n \times d}$ , i.e.,  $V = \{Ax : x \in \mathbb{R}^d\}$ .

A good way to think about the next definition is that we will solve a problem in  $\mathbb{R}^n$  involving an unknown  $d$ -dimensional subspace, by reducing everything to dimension  $s = s(n, d, \varepsilon, \delta)$ . Thus, we want  $s$  (the number of rows in  $S$ ) to be as small as possible.

**Definition:** A random matrix  $S \in \mathbb{R}^{s \times n}$  is called an  $(\varepsilon, \delta, d)$ -*Oblivious Subspace Embedding (OSE)* if

$$\forall A \in \mathbb{R}^{n \times d}, \quad \Pr_S \left[ \forall x \in \mathbb{R}^d, \|SAx\| \in (1 \pm \varepsilon) \|Ax\| \right] \geq 1 - \delta.$$

We next show that it is easy to construct OSE using JLT.

**Exer:** Show that the OSE property is preserved under right-multiplication by a matrix with orthonormal columns, as follows. If  $S \in \mathbb{R}^{s \times n}$  is an  $(\epsilon, \delta, d)$ -OSE matrix, and  $U \in \mathbb{R}^{n \times r}$  is a matrix with orthonormal columns, then  $SU$  is an  $(\epsilon, \delta, \min(r, d))$ -OSE matrix (for the space  $\mathbb{R}^r$ ).

**Theorem 2:** Let  $S \in \mathbb{R}^{s \times n}$  be an  $(\epsilon, \delta, b)$ -JLT. Then  $S$  is also an  $(O(\epsilon), \delta, \epsilon \ln b)$ -OSE.

Remark: For this theorem to produce OSE for dimension  $d$ , we should set  $d = \epsilon \ln b$ , i.e.,  $b = e^{d/\epsilon}$ , which we can achieve using a Gaussian matrix with  $s = O(\epsilon^{-2} \log(b/\delta)) = O(\epsilon^{-2}(\frac{d}{\epsilon} + \log \frac{1}{\delta}))$  rows. A direct construction with sparse columns (and thus fast matrix-vector multiplication) was recently shown by [Cohen, 2016].

Remark 2: In a discussion after the class, we realized that the final inequality we used to bound  $|N|$  is lossy. The proof works even if we set  $b = (1 + 2/\gamma)^d = (1 + 1/\epsilon)^d$  (put differently, it improves to  $d = O(\frac{\ln b}{\ln(1/\epsilon)})$ ), which we can achieve using a Gaussian matrix with  $s = O(\epsilon^{-2} \log(b/\delta)) = O(\epsilon^{-2}(d \log \frac{1}{\epsilon} + \log \frac{1}{\delta}))$  rows.

**Proof:** Was seen in class. The main idea is to use the JLT guarantee on a  $2\epsilon$ -net  $N$  of the unit sphere in  $\mathbb{R}^d$ , then represent arbitrary  $x \in \mathbb{R}^d$  as an infinite (but converging) sum  $x = \sum_{i=0}^{\infty} x_i$ , where each  $x_i$  is a scalar multiple of a net points, and finally use the triangle inequality.

Remark: It is possible to get a better bound by employing a  $1/2$ -net (instead of  $\epsilon$ -net) and expanding  $\|SAx\|^2$  including cross terms.

## 4 Least Squares Regression

**Problem definition:** In *Least Squares Regression*, the input is a matrix  $A \in \mathbb{R}^{n \times d}$  and a vector  $b \in \mathbb{R}^n$ , and the goal is to find  $\operatorname{argmin}\{\|Ax^* - b\| : x^* \in \mathbb{R}^d\}$ .

Informally, when solving a system  $Ax^* = b$  that is over-constrained ( $n \gg d$ ), we do not expect to find an exact solution, and we want to minimize the sum of squared errors  $\sum_i (A_i x^* - b_i)^2$ .

We shall consider  $(1 + \epsilon)$ -approximation, i.e., finding  $x' \in \mathbb{R}^d$  such that

$$\|Ax' - b\| \leq (1 + \epsilon) \min_{x^* \in \mathbb{R}^d} \|Ax^* - b\|. \quad (1)$$

**Theorem 3:** Let  $S \in \mathbb{R}^{n \times s}$  be an  $(\epsilon, \delta, d + 1)$ -OSE matrix. Then for every regression instance  $A \in \mathbb{R}^{n \times d}$  and  $b \in \mathbb{R}^n$ , with high probability, an optimal solution to the regression instance  $\langle SA, Sb \rangle$  is a  $(1 + O(\epsilon))$ -approximation to the instance  $\langle A, b \rangle$ , i.e., such  $x'$  satisfies (1).

This theorem essentially reduces the a regression problem with  $n$  constraints to regression with  $s$  constraints, but we should take into account also the time to compute  $SA$ .

**Proof:** As explained in class, it follows from applying the OSE guarantee on the linear subspace spanned by the columns of  $A$  and by  $b$  (total of  $d + 1$  vectors), and then

$$\|SAx' - Sb\| = \min_{x \in \mathbb{R}^d} \|SAx - Sb\| \leq \frac{1 + \epsilon}{1 - \epsilon} \cdot \min_{x^* \in \mathbb{R}^d} \|Ax^* - b\|.$$