

# On Testing Asymmetry in the Bounded Degree Graph Model

Oded Goldreich\*

January 30, 2021

## Abstract

We consider the problem of testing asymmetry in the bounded-degree graph model, where a graph is called asymmetric if the identity permutation is its only automorphism. Seeking to determine the query complexity of this testing problem, we provide partial results. Considering the special case of  $n$ -vertex graphs with connected components of size at most  $s(n) = \Omega(\log n)$ , we show that the query complexity of  $\epsilon$ -testing asymmetry (in this case) is at most  $O(\sqrt{n} \cdot s(n)/\epsilon)$  and at least  $\Omega(\sqrt{n^{1-O(\epsilon)}/s(n)})$ . In particular, the query complexity of  $o(1/s(n))$ -testing asymmetry is at least  $\Omega(\sqrt{n/s(n)})$ .

In addition, we show that testing asymmetry in the dense graph model is almost trivial.

## Contents

1	Introduction	1
2	In the bounded-degree graph model	2
3	In the dense graph model	7

## 1 Introduction

Property testing refers to probabilistic algorithms of sub-linear complexity for deciding whether a given object has a predetermined property or is far from any object having this property. Such algorithms, called testers, obtain local views of the object by *performing queries* and their performance guarantees are stated with respect to a distance measure that (combined with a distance parameter) determines which objects are considered far from the property.

In the last couple of decades, the area of property testing has attracted significant attention (see, e.g., [5]). Much of this attention was devoted to testing graph properties in a variety of models including the dense graph model [7], and the bounded-degree graph model [8] (surveyed in [5, Chap. 8] and [5, Chap. 9], resp.). We mention, without elaboration, that the known results concerning these models include both results regarding general classes of graph properties and results regarding many natural graph properties. Yet, one natural property that (to the best of our knowledge) was not considered before is *asymmetry*.

A graph is called *asymmetric* if the identity permutation is its only automorphism. Recall that, for a (labeled) graph  $G = (V, E)$  and a bijection  $\phi : V \rightarrow V'$ , we denote by  $\phi(G)$  the graph

---

\*Department of Computer Science, Weizmann Institute of Science, Rehovot, ISRAEL. E-mail: [oded.goldreich@weizmann.ac.il](mailto:oded.goldreich@weizmann.ac.il). Partially supported by the Israel Science Foundation (grant No. 1041/18).

$G' = (V', E')$  such that  $E' = \{\{\phi(u), \phi(v)\} : \{u, v\} \in E\}$ , and say that  $G'$  is isomorphic to  $G$ . The set of automorphisms of the graph  $G = (V, E)$ , denoted  $\text{aut}(G)$ , is the set of permutations that preserve the graph  $G$ ; that is,  $\pi \in \text{aut}(G)$  if and only if  $\pi(G) = G$ .

**Definition 1** (asymmetric and symmetric graphs): *A graph is called asymmetric if its sets of automorphisms is a singleton, which consists of the trivial automorphism (i.e., the identity permutation). Otherwise, the graph is called symmetric.*

It turns out that testing asymmetry in the dense graph model is quite trivial, because, under the corresponding distance measure, every graph is close to being asymmetric (see Section 3). Our focus is on the bounded-degree graph model, where we essentially determine the query complexity for the special case in which all connected components in the graph are small (i.e., have poly-logarithmic size). In other words, we prove.

**Theorem 2** (testing asymmetric graphs with small connected components (in the bounded-degree graph model)): *The query complexity of  $\epsilon$ -testing whether an  $n$ -vertex graph is asymmetric and has connected components of size  $\text{poly}(\log n)$  is at most  $\tilde{O}(n^{0.5}/\epsilon)$  and at least  $\tilde{\Omega}(n^{0.5-O(\epsilon)})$ . In particular, the query complexity of  $o(1/\log n)$ -testing asymmetry is at least  $\tilde{\Omega}(n^{0.5})$ . Furthermore, the upper bound holds for one-sided error testers, whereas the lower bound holds also for general (i.e., two-sided error) testers.*

The results generalize to graphs with connected components of size at most  $s(n) = \Omega(\log n)$ , but in that case the gap between the upper and lower bounds is  $\text{poly}(s(n))$ . Note that, for  $s(n) = o((\log n)/\log \log n)$ , the testing problem is trivial, since the number of bounded-degree  $s$ -vertex graphs is smaller than  $\exp(O(s \log s))$ .<sup>1</sup>

Hence, our state of knowledge is analogous to the state of knowledge regarding (both versions of) the graph isomorphism testing problem.<sup>2</sup> For all (three) problems, we essentially know the query complexity for the special case in which all connected components in the graph are small (i.e., have poly-logarithmic size), but we are otherwise ignorant about the general case: In particular, *do testers of sublinear query complexity exist for these problems (in the general case)? Or, on the other hand, is the general case harder than the special case?*

## 2 In the bounded-degree graph model

In the bounded-degree model, graphs are represented by their incidence functions and distances are measured as the ratio of the number of differing incidences over the maximal number of edges. Specifically, for a degree bound  $d \in \mathbb{N}$ , we represent a graph  $G = ([n], E)$  of maximum degree  $d$  by the incidence function  $g : [n] \times [d] \rightarrow [n] \cup \{0\}$  such that  $g(v, i)$  indicates the  $i^{\text{th}}$  neighbor of  $v$

<sup>1</sup>This implies that an  $n$ -vertex graph that consists of connected components of size at most  $s(n) = o((\log n)/\log \log n)$  must have a few identical components, and is thus symmetric.

<sup>2</sup>For testing Graph Isomorphism in the bounded-degree graph model, the following is known [6].

1. The query complexity of testing isomorphism to a fixed  $n$ -vertex graph is  $\tilde{\Omega}(n^{1/2})$ .
2. The query complexity of testing isomorphism between two  $n$ -vertex graphs is  $\tilde{\Omega}(n^{2/3})$ .

The lower bounds are shown by using graphs that have connected components of size  $\text{poly}(\log n)$ , and in this case the lower bounds are tight [6]. We mention that, unlike Theorem 2, one-sided error testing of isomorphism (even to a fixed graph) has linear query complexity [6, Thm. 2.5].

(where  $g(v, i) = 0$  indicates that  $v$  has less than  $i$  neighbors). The distance between the graphs  $G = ([n], E)$  and  $G' = ([n], E')$  is defined as the symmetric difference between  $E$  and  $E'$  over  $dn/2$ , and oracle access to a graph means oracle access to its incidence function.

**Definition 3** (testing graph properties in the bounded-degree graph model): *For a fixed degree bound  $d$ , a tester for a graph property  $\Pi$  is a probabilistic oracle machine that, on input parameters  $n$  and  $\epsilon$ , and oracle access to (the incidence function of) an  $n$ -vertex graph  $G = ([n], E)$  of maximum degree  $d$ , outputs a binary verdict that satisfies the following two conditions.*

1. *If  $G \in \Pi$ , then the tester accepts with probability at least  $2/3$ .*
2. *If  $G$  is  $\epsilon$ -far from  $\Pi$ , then the tester accepts with probability at most  $1/3$ , where  $G$  is  $\epsilon$ -far from  $\Pi$  if for every  $n$ -vertex graph  $G' = ([n], E') \in \Pi$  of maximum degree  $d$  it holds that the symmetric difference between  $E$  and  $E'$  has cardinality that is greater than  $\epsilon \cdot dn/2$ .*

*If the tester accepts every graph in  $\Pi$  with probability 1, then we say that it has one-sided error; otherwise, we say that it has two-sided error.*

(Throughout this work, we consider undirected simple graphs (i.e., no self-loops and parallel edges).)

The query complexity of a tester for  $\Pi$  is a function (of the parameters  $d, n$  and  $\epsilon$ ) that represents the number of queries made by the tester on the worst-case  $n$ -vertex graph of maximum degree  $d$ , when given the proximity parameter  $\epsilon$ . Fixing  $d$ , we typically ignore its effect on the complexity (equiv., treat  $d$  as a hidden constant). The query complexity of  $\epsilon(n)$ -testing  $\Pi$  is defined as the query complexity of testing when the proximity parameter is set to  $\epsilon(n)$ ; that is, we say that the query complexity of  $\epsilon(n)$ -testing  $\Pi$  is at least  $Q(n)$  if distinguishing between  $n$ -vertex graphs in  $\Pi$  and  $n$ -vertex graphs that are  $\epsilon(n)$ -far from  $\Pi$  requires at least  $Q(n)$  queries.

**Theorem 4** (Theorem 2, restated): *For  $s : \mathbb{N} \rightarrow \mathbb{N}$ , let  $\Pi^{(s)} = \bigcup_{n \in \mathbb{N}} \Pi_n^{(s)}$  such that  $\Pi_n^{(s)}$  is the set of asymmetric  $n$ -vertex graphs that have connected components of size at most  $s(n)$ . Then, for every degree bound  $d \geq 3$ , the following holds.*

1. *If  $s(n) = \Omega((\log n)/\log \log n)$ , then the query complexity of  $\epsilon$ -testing  $\Pi_n^{(s)}$  (in the bounded-degree graph model) is  $\Omega((n/s(n))^{0.5-O(\epsilon)})$ .  
In particular, the query complexity of  $(1/(3d \cdot s(n)))$ -testing  $\Pi_n^{(s)}$  is  $\Omega((n/s(n))^{0.5})$ .*
2. *There exists a one-sided error  $\epsilon$ -tester for  $\Pi^{(s)}$  (in the bounded-degree graph model) that makes  $O(n^{0.5} \cdot s(n)/\epsilon)$  queries, and runs in time  $\tilde{O}(n^{0.5}/\epsilon) \cdot \text{poly}(s(n))$ .*

We stress that Part 1 holds also for two-sided error testers. Recall that, for  $s(n) = o((\log n)/\log \log n)$ , the testing problem is trivial, since the number of bounded-degree  $s$ -vertex graphs is smaller than  $\exp(O(s \log s)) < n/s(n)$ . Theorem 4 follows by combining Propositions 6 and 7, which are stated and proved next. Actually, we start by establishing the special case of Part 1 (i.e., the case of  $\epsilon = (1/(3d \cdot s(n)))$ ).

**Proposition 5** (special case of the lower bound): *For every  $d \geq 3$  and any  $s : \mathbb{N} \rightarrow \mathbb{N}$  such that  $s(n) = \Omega((\log n)/\log \log n)$ , the query complexity of  $(1/(3d \cdot s(n)))$ -testing whether an  $n$ -vertex graph is asymmetric is  $\Omega((n/s(n))^{1/2})$ . This holds even if it is guaranteed that the tested graph consists of connected components of size at most  $s(n)$ .*

**Proof:** We use the following facts, proved in [2, 3]: (F1) most  $d$ -regular  $s$ -vertex graphs are asymmetric, and (F2) their number exceeds  $N_d(s) = \Omega(s/d!)^{ds/2}$ . Note that (F1) holds even if we require the graphs to be connected, since most  $d$ -regular graphs are actually expanders. Hence, for some constant  $c$  and  $s(n) = \frac{c \log_2 n}{d \log_2 \log_2 n}$  it holds that  $\frac{N_d(s(n))}{s(n)!} > 2^{(0.5d-1)c \log_2 n - o(\log n)}$ , which is larger than  $n$  when  $c > 2/(d-2)$ . It follows that there exists a collection, denoted  $C$ , of  $m = n/s(n)$  non-isomorphic  $s(n)$ -vertex  $d$ -regular graphs that are asymmetric and connected. The claim of the proposition follows by showing that  $\Omega(\sqrt{m})$  queries are necessary for distinguish the following two distributions:

1. A random isomorphic copy of the  $n$ -vertex graph  $G_1$  that consists of copies of all graphs in  $C$ ; that is,  $G_1$  consists of  $m$  connected components such that each graph in  $C$  appears as a connected component.
2. A random isomorphic copy of the  $n$ -vertex graph that consists of two copies of  $m/2$  random graphs in  $C$ ; that is, we first select a random  $m/2$ -subset of  $C$ , denoted  $C'$ , and take a random isomorphic copy of the  $n$ -vertex graph  $G_{C'}$  that consists of two copies of each graph in  $C'$ .

Note that each graph in the support of the first distribution is asymmetric, whereas each graph in the support of the second distribution is  $(1/(3d \cdot s(n)))$ -far being asymmetric. The latter claim holds because making  $G_{C'}$  asymmetric requires modifying the incidence of at least one vertex in at least  $m/2$  of its connected components, which amounts to at least  $\frac{m}{4} = \frac{n}{4s(n)} > \frac{1}{3d \cdot s(n)} \cdot dn/2$  edge-modifications.

The fact that  $\Omega(\sqrt{m})$  queries are necessary to distinguish the foregoing two distributions is proved by the “birthday” argument. Specifically, when making  $q$  queries to a graph drawn from the second distribution, we encounter vertices in two different connected components that are isomorphic to the same graph (in  $C$ ) with probability at most  $\binom{q}{2}/|C'|$ , where  $|C'| = m/2$ . Whenever this event does not occur, the answers are distributed identically to the way they are distributed when querying a graph drawn from the first distribution. ■

**Proposition 6** (lower bound on testing asymmetric graphs (in the bounded-degree graph model)): *For every  $d \geq 3$  and any  $s : \mathbb{N} \rightarrow \mathbb{N}$  such that  $s(n) = \Omega((\log n)/\log \log n)$ , the query complexity of  $\epsilon$ -testing the set of  $n$ -vertex asymmetric graphs that consist of connected components of size at most  $s$  is  $\Omega((n/s(n))^{0.5-O(\epsilon)})$ .*

**Proof:** We use the same ingredients as in the proof of Proposition 5, but generalize the argument as follows. Specifically, recall that, for  $s(n) = \frac{c \log_2 n}{d \log_2 \log_2 n}$ , we denote by  $C$  a collection of  $m = n/s(n)$  non-isomorphic  $s(n)$ -vertex  $d$ -regular graphs that are asymmetric expanders. Letting  $t = n^{O(\epsilon)}$ , the claim of the proposition follows by showing that  $\Omega(\sqrt{m/t})$  queries are necessary for distinguish the following two distributions:

1. A random isomorphic copy of the  $n$ -vertex graph  $G_1$  that consists of copies of all graphs in  $C$ ; that is,  $G_1$  consists of  $m$  connected components such that each graph in  $C$  appears as a connected component.
2. A random isomorphic copy of the  $n$ -vertex graph that consists of  $t$  copies of  $m/t$  random graphs in  $C$ ; that is, we first select a random  $m/t$ -subset of  $C$ , denoted  $C'$ , and take a random isomorphic copy of the  $n$ -vertex graph  $G_{C'}$  that consists of  $t$  copies of each graph in  $C'$ .

The first distribution is defined exactly as in the proof of Proposition 5, whereas the second distribution in the latter proof corresponds to the special case of  $t = 2$ . Recalling that each graph in the support of the first distribution is asymmetric, the key observation here is that each graph in the support of the second distribution is  $\epsilon$ -far from being asymmetric. We prove this claim in two steps.

First, note that if  $G_{C'}$  is  $\epsilon$ -close to being asymmetric, then it can be made asymmetric by making  $O(\epsilon dn)$  edge modifications, while preserving the partition of its vertices to connected components. This is the case because the cost of splitting a connected component into two parts is only a constant factor cheaper than the cost of modifying its smaller part arbitrarily, since each connected component is an expander. Second, making the graph  $G_{C'}$  asymmetric, while preserving the partition of its vertices to connected components, requires mapping the  $t$  copies of each graph in  $C'$  to  $t$  different  $s(n)$ -vertex graphs. The point is that the number of  $s$ -vertex  $d$ -regular graphs that are  $\epsilon'$ -close to a given graph is smaller than  $\binom{ds}{2\epsilon' \cdot ds} \cdot (s+1)^{2\epsilon' \cdot ds} < (ds)^{4\epsilon' \cdot ds} < n^{4c\epsilon'}$ . Using  $\epsilon' = O(\epsilon)$  and  $n^{4c\epsilon'} = o(t)$ , it follows that making  $G_{C'}$  asymmetric (while preserving its connected components) requires modifying at least  $2\epsilon' \cdot ds$  incidences in almost each of its connected components, which amounts to more than  $m \cdot \epsilon' \cdot ds = \epsilon' \cdot dn$  edge modifications. Using the first step, the claim (that  $G_{C'}$  is  $\epsilon$ -far from being asymmetric) follows.

The fact that  $\Omega(\sqrt{m/t})$  queries are necessary to distinguish the foregoing two distributions is proved by the ‘‘birthday’’ argument (as in the proof of Proposition 5). Specifically, when making  $q$  queries to a graph drawn from the second distribution, we encounter vertices in two different connected components that are isomorphic to the same graph (in  $C$ ) with probability at most  $\binom{q}{2}/|C'|$ , where  $|C'| = m/t$ . ■

**Proposition 7** (upper bound on testing asymmetric graphs (in the bounded-degree graph model)): *For every  $d \geq 3$ , there exists a one-sided error tester of query complexity  $O(n^{1/2} \cdot s/\epsilon)$  for the set of  $n$ -vertex asymmetric graphs that consist of connected components of size at most  $s$ . Furthermore, the running time of the tester is  $\tilde{O}(n^{1/2}/\epsilon) \cdot \text{poly}(s)$ .*

**Proof:** On input parameters  $n, s$  and  $\epsilon > 0$ , and oracle access to a graph  $G = ([n], E)$ , the algorithm proceeds as follows.

1. It selects uniformly at random  $m = O(\sqrt{n}/\epsilon)$  vertices  $v_1, \dots, v_m \in [n]$ .
2. For each  $i \in [m]$ , the algorithm starts a (e.g., BFS) exploration of the connected component in which  $v_i$  resides, and halts rejecting if it discovers a connected component having more than  $s$  vertices.
3. If for some  $i \in [m]$ , the connected component explored from  $v_i$  is symmetric, then the algorithm halts rejecting.
4. If for some  $i, j \in [m]$ , the connected components explored from  $v_i$  and  $v_j$  are different but isomorphic (i.e.,  $v_i$  does not reside in the same connected component as  $v_j$  but these two connected components are isomorphic), then the algorithm halts rejecting.

If the algorithm did not reject, then it accepts.

The query complexity of this algorithm is  $O(m \cdot s)$ , while its running time is dominated by Steps 3 and 4. Observe, however, that Steps 3 and 4 can be implemented in time  $\tilde{O}(m) \cdot \text{poly}(s)$

by using the canonical labeling algorithm (for bounded-degree graphs) of [1] (along with a sorting algorithm).

Let  $\Pi = \Pi_n^{(s)}$  denote the set of  $n$ -vertex asymmetric graphs that consist of connected components of size at most  $s$ . Evidently, the algorithm accepts each graph in  $\Pi$  with probability 1. On the other hand, if  $G$  is  $\epsilon$ -far from  $\Pi$ , then one of the following three cases must hold.

**Case 1:** At least  $\epsilon n/6$  of its vertices reside in connected components of size greater than  $s$ .

In this case, Step 2 of the algorithm rejects (w.h.p.).

**Case 2:** At least  $\epsilon n/6$  of its vertices reside in connected components of size at most  $s$  that are symmetric.

In this case, Step 3 of the algorithm rejects (w.h.p.).

**Case 3:** At most  $\epsilon n/3$  of its vertices reside in connected components that are either of size exceeding  $s$  or are symmetric.

In this case, let  $S$  denote the set of all other vertices (i.e., vertices that reside in asymmetric connected components of size at most  $s$ ), and let  $G_S$  denote the subgraph of  $G$  induced by  $S$ .

Consider the graph  $G'$  that results by augmenting  $G_S$  with  $(n - |S|)/s$  connected components (each of size  $s$ ) that are neither symmetric nor isomorphic to any other connected component (where the existence of such a collection of  $s$ -vertex graphs has been established in the first paragraph of the proof of Proposition 5). Recalling that  $G$  is  $\epsilon$ -far from  $\Pi$  and  $n - |S| \leq \epsilon n/3$ , it follows that  $G'$  is  $\epsilon/3$ -far from being asymmetric.

Let  $C_1, \dots, C_{m'}$  denote the connected components of  $G_S$ , and recall that each  $C_i$  has at most  $s$  vertices. Consider the equivalence relation, denoted  $\equiv$ , defined by graph isomorphism (over the set of  $C_i$ 's); that is,  $C_i \equiv C_j$  if and only if  $C_i$  is isomorphic to  $C_j$ . Let  $n_k$  denote the number of  $k$ -vertex connected components that reside in equivalence classes that has more than a single  $C_i$ ; that is,

$$n_k = |\{i \in [m'] : |C_i| = k \ \& \ \exists j \neq i \text{ s.t. } C_i \equiv C_j\}|$$

where  $|C_i|$  denotes the number of vertices in  $C_i$ . Then,  $\sum_{k \in [s]} n_k \cdot k \geq \epsilon n/6$  must hold, because otherwise  $G'$  is  $\epsilon/3$ -close to being asymmetric; to see this, replace the connected components in the *non-singleton equivalence classes* by asymmetric connected components of size  $s$  that are not isomorphic to any other connected component (see the foregoing comment regarding the existence of such a collection).

Now, if we take a sample of  $\Theta(\epsilon^{-1}\sqrt{n})$  vertices, then it is very likely that  $\Theta(\sqrt{n})$  of these vertices hit connected components in the non-singleton equivalence classes. Recalling that  $\sum_{k \in [s]} n_k \cdot k \leq n$ , we infer that this sample is likely to hit two different elements of the same class. This holds because, with high probability, we are likely to have several classes hit by at least two samples, and with probability at least  $1/2$  each of these pairs of samples hit different  $C_i$ 's in the relevant class.

Hence, in each of these cases, the algorithm rejects with high probability, which establishes our claim. ■

**On testing the set of symmetric graphs.** We mention that testing the set of symmetric graphs is almost trivial; specifically, the query complexity is 0 if  $\epsilon \geq 4/n$ , and  $dn = O(d/\epsilon)$  otherwise. This is the case because, with respect to a degree bound  $d$ , every  $n$ -vertex graph is  $\frac{2d}{dn/2}$ -close to being symmetric (e.g., by making two vertices isolated).

### 3 In the dense graph model

In the dense graph model, a graph  $G = ([n], E)$  is represented by its adjacency predicate,  $g : [n] \times [n] \rightarrow \{0, 1\}$ , such that  $g(u, v) = 1$  if and only if  $\{u, v\} \in E$ . The distance between the graphs  $G = ([n], E)$  and  $G' = ([n], E')$  is defined as the symmetric difference between  $E$  and  $E'$  over  $\binom{[n]}{2}$ , and oracle access to a graph means oracle access to its adjacency predicate.

**Definition 8** (testing graph properties in the dense graph model): *A tester for a graph property  $\Pi$  is a probabilistic oracle machine that, on input parameters  $n$  and  $\epsilon$ , and oracle access to (the adjacency predicate of) an  $n$ -vertex graph  $G = ([n], E)$ , outputs a binary verdict that satisfies the following two conditions.*

1. *If  $G \in \Pi$ , then the tester accepts with probability at least  $2/3$ .*
2. *If  $G$  is  $\epsilon$ -far from  $\Pi$ , then the tester accepts with probability at most  $1/3$ , where  $G$  is  $\epsilon$ -far from  $\Pi$  if for every  $n$ -vertex graph  $G' = ([n], E') \in \Pi$  it holds that the symmetric difference between  $E$  and  $E'$  has cardinality that is greater than  $\epsilon \cdot \binom{[n]}{2}$ .*

The query complexity of a tester for  $\Pi$  is a function (of the parameters  $n$  and  $\epsilon$ ) that represents the number of queries made by the tester on the worst-case  $n$ -vertex graph, when given the proximity parameter  $\epsilon$ . In this section, we show that testing the set of asymmetric graphs in the dense graph model is almost trivial; specifically, the query complexity is 0 if  $\epsilon > O((\log n)/n)$ , and  $\binom{[n]}{2} = \tilde{O}(1/\epsilon^2)$  otherwise. This holds because in the first case (i.e.,  $\epsilon > O((\log n)/n)$ ), all  $n$ -vertex graphs are  $\epsilon$ -close to being asymmetric (see Proposition 9), whereas in the second case one can afford to retrieve the entire graph.

**Proposition 9** (all graphs are close to being asymmetric): *In the dense graph model, every  $n$ -vertex graph  $G$  is  $\frac{O(\log n)}{n}$ -close to being asymmetric.*

**Proof:** Given an arbitrary graph  $G = ([n], E)$ , we construct a random variant of it, denoted  $G'$ , by re-randomizing  $O(n \log n)$  of its adjacencies, and show that (w.h.p.) the resulting graph is asymmetric. Specifically, we consider the following “randomized” version.

**Construction 9.1** (construction of  $G'$ ): *Given an arbitrary graph  $G = ([n], E)$ , we proceed as follows.*

1. *Select an arbitrary subset,  $S$ , of  $\ell = O(\log n)$  vertices in  $G$ .*
2. *Replace the subgraph of  $G$  induced by  $S$  with a random  $\ell$ -vertex graph.*
3. *Replace the bipartite subgraph that connects  $S$  and  $[n] \setminus S$  by a random bipartite graph; that is, for each  $s \in S$  and  $v \in [n] \setminus S$ , the edge  $\{s, v\}$  is contained in the resulting graph  $G'$  with probability  $1/2$ .*

We shall first show that, with very high probability, the subgraph of  $G'$  induced by  $S$  is not isomorphic to the subgraph of  $G'$  that is induced by any other  $\ell$ -subset.

**Claim 9.2** (uniqueness of  $S$ ): *For every  $\ell$ -subset  $S$  fixed in Step 1 of Construction 9.1, with high probability over Steps 2 and 3, for every  $\ell$ -subset  $S' \neq S$  of  $[n]$ , the subgraph of  $G'$  induced by  $S'$  is not isomorphic to the subgraph of  $G'$  induced by  $S$ .*

**Proof:** The case of  $S' \cap S = \emptyset$  is easy, because in this case the subgraph of  $G'$  induced by  $S'$  is fixed in Step 1 (since it equals the subgraph of  $G$  induced by  $S'$ ), whereas a random  $\ell$ -vertex graph (as selected in Step 2) is isomorphic to this fixed graph with probability at most  $(\ell!) \cdot 2^{-\binom{\ell}{2}} \ll \binom{n}{\ell}^{-1}$ , where the inequality uses a sufficiently large  $\ell = O(\log n)$ . Hence, we can afford to take a union bound over all  $\ell$ -subsets that are disjoint of  $S$ . However, for sets that are not disjoint of  $S$ , the foregoing probability bound does not hold, and a more careful analysis is called for. Nevertheless, the foregoing analysis does provide a good warm-up towards the rest.

First, for each  $\ell$ -set  $S' \subset [n]$  such that  $S' \neq S$ , we shall upper-bound the probability that the subgraphs of  $G'$  induced by  $S$  and  $S'$  are isomorphic as a function of  $|S \cap S'|$ . For every bijection  $\pi : S \rightarrow S'$ , let  $\text{FP}(\pi) \stackrel{\text{def}}{=} \{v \in S : \pi(v) = v\}$  denote the set of fixed-points of  $\pi$ , and note that  $|\text{FP}(\pi)| \leq \ell - 1$  (since  $S \neq S'$ ). Now, letting  $G_R$  denote the subgraph of  $G$  induced by  $R$ , we claim that the probability that there exists a bijection  $\pi : S \rightarrow S'$  such that  $\pi(G'_S) = G'_{S'}$  is upper-bounded by

$$\sum_{\pi: S \xrightarrow{1-1} S'} \min \left( 2^{-|\text{FP}(\pi)| \cdot (\ell - |\text{FP}(\pi)|) / 3}, 2^{-\binom{\ell - |\text{FP}(\pi)|}{2} / 3} \right) \quad (1)$$

$$\begin{aligned} &\leq \sum_{f \in \{0, \dots, |S \cap S'|\}} \frac{\ell!}{f!} \cdot 2^{-\max(6 \cdot f \cdot (\ell - f), (\ell - f) \cdot (\ell - f - 1)) / 18} \\ &< \frac{\ell!}{|S \cap S'|!} \cdot 2^{-\Omega((\ell - |S \cap S'|) \cdot \ell)} \end{aligned} \quad (2)$$

where  $f$  represents the size of  $\text{FP}(\pi)$ . To justify the upper bound claimed in Eq. (1), consider an arbitrary bijection  $\pi : S \rightarrow S'$ , and identify a set  $I \subseteq S \setminus \text{FP}(\pi)$  such that  $\pi(I) \cap I = \emptyset$  and  $|I| \geq (\ell - |\text{FP}(\pi)|) / 3$ . Letting  $e_{G'}(u, v) = 1$  if  $\{u, v\}$  is an edge in  $G'$  and  $e_{G'}(u, v) = 0$  otherwise, observe that  $\pi(G'_S) = G'_{S'}$  if and only if  $e_{\pi(G')}(u, v) = e_{G'}(\pi(u), \pi(v))$  for every  $\{u, v\} \in \binom{S}{2}$ . Noting that  $e_{\pi(G')}(u, v) = e_{G'}(u, v)$ , the first bound in Eq. (1) is justified by

$$\begin{aligned} &\Pr_{G'} \left[ \forall (u, v) \in \binom{S}{2} : e_{\pi(G')}(u, v) = e_{G'}(\pi(u), \pi(v)) \right] \\ &\leq \Pr_{G'} [\forall (u, v) \in \text{FP}(\pi) \times I : e_{G'}(u, v) = e_{G'}(\pi(u), \pi(v))] \\ &= \prod_{(u, v) \in \text{FP}(\pi) \times I} \Pr_{G'} [e_{G'}(u, v) = e_{G'}(\pi(u), \pi(v))] \\ &= 2^{-|\text{FP}(\pi)| \cdot |I|} \\ &\leq 2^{-|\text{FP}(\pi)| \cdot (\ell - |\text{FP}(\pi)|) / 3} \end{aligned}$$

where the equalities are due to the disjointness of the sets  $\text{FP}(\pi) \times I$  and  $\text{FP}(\pi) \times \pi(I)$  (to the fact that  $\pi(u) = u$  for every  $u \in \text{FP}(\pi)$ ), and to the fact that the incidences of all vertices in  $\text{FP}(\pi) \subseteq S$  are random. Similarly, we justify the second bound in Eq. (1) by



$$\begin{aligned}
& \Pr_{G'} \left[ \forall \{u, v\} \in \binom{S}{2} : e_{\pi(G')}(\pi(u), \pi(v)) = e_{G'}(\pi(u), \pi(v)) \right] \\
& \leq \Pr_{G'} \left[ \forall \{u, v\} \in \binom{I}{2} : e_{G'}(u, v) = e_{G'}(\pi(u), \pi(v)) \right] \\
& = \prod_{\{u, v\} \in \binom{I}{2}} \Pr_{G'} [e_{G'}(u, v) = e_{G'}(\pi(u), \pi(v))] \\
& = 2^{-\binom{|I|}{2}} \\
& \leq 2^{-\binom{\ell - |\text{FP}(\pi)|/3}{2}}
\end{aligned}$$

where the equalities are due to the disjointness of the sets  $\binom{I}{2}$  and  $\binom{\pi(I)}{2}$ , and to the fact that the incidences of all vertices in  $I \subseteq S \setminus \text{FP}(\pi) \subseteq S$  are random.

Combining Eq. (1)&(2) with a union bound over all  $\ell$ -subsets  $S' \subset [n]$  that are different from  $S$ , we upper-bound the probability that the subgraphs of  $G'$  induced by  $S$  and by some other  $\ell$ -set are isomorphic by

$$\sum_{S' \in \binom{[n]}{\ell} \setminus \{S\}} \frac{\ell!}{|S \cap S'|!} \cdot 2^{-\Omega((\ell - |S \cap S'|) \cdot \ell)} = \sum_{i \in \{0, \dots, \ell-1\}} \binom{\ell}{i} \cdot \binom{n-i}{\ell-i} \cdot \frac{\ell!}{i!} \cdot 2^{-\Omega((\ell-i) \cdot \ell)} \quad (3)$$

where the index  $i$  represents the size of the intersection with  $S$ . Using a sufficiently large  $\ell = O(\log n)$ , we have

$$\begin{aligned}
\sum_{i \in \{0, \dots, \ell-1\}} \binom{\ell}{i} \cdot \binom{n-i}{\ell-i} \cdot \frac{\ell!}{i!} \cdot 2^{-\Omega((\ell-i) \cdot \ell)} &= \sum_{i \in \{0, \dots, \ell-1\}} \binom{\ell}{i}^2 \cdot \binom{n-i}{\ell-i} \cdot \frac{(n-i)!}{(n-\ell)!} \cdot 2^{-\Omega((\ell-i) \cdot \ell)} \\
&< \sum_{i \in \{0, \dots, \ell-1\}} n^{\ell-i} \cdot \binom{\ell}{i}^2 \cdot 2^{-\Omega((\ell-i) \cdot \ell)} \\
&< \ell \cdot \max_{i \in \{0, \dots, \ell-1\}} \left\{ n^{\ell-i} \cdot \binom{\ell}{i}^2 \cdot 2^{-\Omega((\ell-i) \cdot \ell)} \right\} \\
&= \ell \cdot (n \cdot \ell^2 \cdot 2^{-\Omega(\ell)})
\end{aligned}$$

which is  $o(1)$ . The claim follows. ■

**Conclusion.** Using Claim 9.2, we claim that (w.h.p.) the graph  $G'$  is asymmetric. This holds because each of the following claims holds with high probability.

1. Any automorphism of the graph  $G'$  maps the set  $S$  to itself.  
(Indeed, this is due to Claim 9.2.)
2. The subgraph of  $G'$  induced by  $S$  is asymmetric.  
(Recall that by [4], almost all  $\ell$ -vertex graphs are asymmetric.)

3. Any vertex  $v \in [n] \setminus S$  has a different “neighborhood pattern” with respect to  $S$ ; that is, for every  $u \neq v \in [n] \setminus S$ , there exists  $w \in S$  such that  $\{u, w\}$  is an edge in  $G'$  if and only if  $\{v, w\}$  is not an edge in  $G'$ .

By combining Conditions 1 and 2, it follows that any automorphism of the graph  $G'$  maps each vertex  $w \in S$  to itself, whereas by Condition 3 such an isomorphism must map each  $v \in [n] \setminus S$  to itself. Hence, the claim (that  $G'$  is asymmetric) follows, and the proposition follows by noting that  $G'$  is  $\frac{\ell \cdot n}{n^2}$ -close to  $G$ . ■

**On testing the set of symmetric graphs.** We mention that testing the set of symmetric graphs is also almost-trivial; specifically, the query complexity is 0 if  $\epsilon \geq 1/n$ , and  $\binom{n}{2} = O(1/\epsilon^2)$  otherwise. This is the case because each  $n$ -vertex graph is  $\frac{1}{n}$ -close to being symmetric, since by [4, Thm. 1] any  $n$ -vertex graph can be made symmetric by modifying the edge relation of at most  $\frac{n-1}{2}$  vertex-pairs. (Note that an upper bound of  $n - 1$  is obvious by picking two vertices  $u$  and  $v$ , and modifying the neighborhood of  $u$  to equal that of  $v$ .)

## References

- [1] L. Babai and E.M. Luks. Canonical Labeling of Graphs. In *15th ACM Symposium on the Theory of Computing*, pages 171–183, 1983.
- [2] B. Bollobas. Distinguishing Vertices of Random Graphs. *North-Holland Mathematics Studies*, Vol. 62, pages 33–49, 1982.
- [3] B. Bollobas. The Asymptotic Number of Unlabelled Regular Graphs. *J. Lond. Math. Soc.*, Vol. 26, pages 201–206, 1982.
- [4] P. Erdos and A. Renyi. Asymmetric Graphs. *Acta Mathematica Hungarica*, Vol. 14 (3), pages 295–315, 1963.
- [5] O. Goldreich. *Introduction to Property Testing*. Cambridge University Press, 2017.
- [6] O. Goldreich. Testing Isomorphism in the Bounded-Degree Graph Model. *ECCC*, TR19-102, 2019.
- [7] O. Goldreich, S. Goldwasser, and D. Ron. Property testing and its connection to learning and approximation. *Journal of the ACM*, pages 653–750, July 1998. Extended abstract in *37th FOCS*, 1996.
- [8] O. Goldreich and D. Ron. Property Testing in Bounded Degree Graphs. *Algorithmica*, Vol. 32 (2), pages 302–343, 2002.